

Energy-preserving methods and B-series

Elena Celledoni

joint work with

D. McLaren, R.I. McLachlan, B. Owren, G.R. Quispel, W. Wright

Preliminary abstract

Given the system of ordinary differential equations (ODEs)

$$\dot{y} = f(y), \quad y(0) = y_0 \in \mathbf{R}^n,$$

assume $H : \mathbf{R}^n \rightarrow \mathbf{R}$ is an invariant or the energy of the system, and that we have $f(y) = S \nabla H(y)$ with S a $n \times n$, skew-symmetric matrix¹.

To derive energy-preserving numerical integration methods for these problems one can use discrete gradient techniques, [3], [4]. Such methods rely on appropriate approximations of $\nabla H(y)$ and S (in the case S depends on y), and in general can not be expanded in a B-series. One surprising exception is the average vector field (AVF) method, namely

$$y_{n+1} - y_n = h \int_0^1 f((1 - \xi)y_n + \xi y_{n+1}) d\xi, \quad y_n \approx y(t_n), \quad n = 0, 1, \dots \quad (1)$$

Under the assumption that S is a constant matrix, this is both a B-series method and a discrete gradient method, as recently observed by Quispel and McLaren [5].

Faou, Hairer and Pham, [2], and Chartier, Faou and Murua, [1], characterized precisely all energy preserving B-series for Hamiltonian problems ($\dot{y} = S \nabla H(y)$, with S the canonical symplectic matrix).

The recent work on (1) brings new possibilities for the construction of concrete energy-preserving, B-series methods.

We will discuss possible extensions of (1) within the class of B-series methods. To illustrate the performance of the methods some numerical experiments on partial differential equations will be presented.

¹This implies that the energy H is preserved along the solution of the ODE, that is

$$\frac{dH(y(t))}{dt} = \nabla H(y)^T \cdot f(y) = \nabla H(y)^T \cdot S \nabla H(y) = 0.$$

References

- [1] P. Chartier, E. Faou, and A. Murua. An algebraic approach to invariant preserving integrators: the case of quadratic and Hamiltonian invariants. *Numer. Math.*, 103(4):575–590, 2006.
- [2] E. Faou, E. Hairer, and T. L. Pham. Energy conservation with non-symplectic methods: examples and counter-examples. *BIT*, 44(4):699–709, 2004.
- [3] O. Gonzalez. Time integration and discrete Hamiltonian systems. *J. Nonlinear Sci.*, 6(5):449–467, 1996.
- [4] R. I. McLachlan, G. R. W. Quispel, and N. Robidoux. Geometric integration using discrete gradients. *R. Soc. Lond. Philos. Trans. Ser. A Math. Phys. Eng. Sci.*, 357(1754):1021–1045, 1999.
- [5] G. R. W. Quispel and D. I. McLaren. A new class of energy-preserving numerical integration methods. *Journal of Phys. A: Math. Theor.*, 41:xxx–xxx, 2008. To appear.