



MA2501 Numeriske Metoder  
Olivier Verdier

## Training Assignment 1

2012-01-12

This assignment has 6 tasks.

**Exercise 1.** We are looking for a solution of the equation  $x^3 + x - 2 = 0$ .

- 1.a) Write down one step of Newton's method.
- 1.b) Compute the result after two steps, starting from the initial guess  $x_0 = 0$ .
- 1.c) Write down one step of Newton's method as an expression computable in a computer, as was done in the lecture.

$$x = x - (x^{**3} + x - 2)/(3*x + 1)$$

**Exercise 2.** How many steps are necessary for Newton's method to converge when computing the root of  $f(x) = ax + b$ ? Does it depend on the starting point?

**Exercise 3.** Write down Newton's method for the function  $f(x) = \frac{1}{x}$ . What happens? Calculate the result after 50 iterations, assuming that the starting value is 1.

**Exercise 4.** The sequence  $u_n$  is defined implicitly, for a given function  $f$  and a small, positive real number  $h$ , as

$$\frac{u_{n+1} - u_n}{h} = f(u_{n+1}).$$

- 4.a) If you also fix  $u_n$ , find a function of which  $u_{n+1}$  is the root.
- 4.b) Write down one step of Newton's method, that should converge to that root.
- 4.c) What initial condition would you choose to start the iteration?

**Exercise 5.** A continuous functions which changes its sign in an interval  $[a, b]$ , i.e.  $f(a)f(b) < 0$ , has at least one zero, or *root*, in this interval. In other words, there exists at least one point  $r \in [a, b]$  such that

$$f(r) = 0.$$

Such a root  $r$  can be found by the *bisection method*, which we describe now.

This method starts from the given interval  $[a, b]$ . Then it investigates the sign changes in the subintervals  $[a, \frac{a+b}{2}]$  and  $[\frac{a+b}{2}, b]$ . If the sign changes in the first subinterval  $b$  is redefined to be

$$b := \frac{a + b}{2}$$

otherwise,  $a$  is redefined in the same manner to

$$a := \frac{a + b}{2},$$

and the process is repeated until the quantity  $|b - a|$  is less than a given tolerance.

Show that the bisection method converges, if the assumption are fulfilled.

**Exercise 6.** Assuming that Newton's method and bisection method converge, one can show the following estimates of their respective errors, *provided that the initial error  $e_0$  is sufficiently small*:

$$\begin{array}{ll} |e_{n+1}^N| \leq C|e_n^N|^2 & \text{Newton} \\ |e_{n+1}^B| \leq C' \frac{|e_n^B|}{2} & \text{Bisection} \end{array}$$

Which method, in general, will be fastest to converge?