



Faglig kontakt under eksamen:
Idun Reiten (telefon: 73 53 45 79, 99 24 45 39)

MNFMA318, Rings and modules
English

Saturday, Desember 7, 2002

Time: 9-13

Permitted aids: None

Grades to be announced: Monday, January 6, 2003

Problem 1

Let \mathbb{R} denote the field of real numbers and let

$$R = \left\{ \begin{pmatrix} a & 0 & 0 & 0 \\ b & c & 0 & 0 \\ d & 0 & e & 0 \\ f & 0 & 0 & g \end{pmatrix} ; a, b, c, d, e, f, g \in \mathbb{R} \right\} \quad \text{and} \quad I = \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 \\ b & 0 & 0 & 0 \\ d & 0 & 0 & 0 \\ f & 0 & 0 & 0 \end{pmatrix} ; b, d, f \in \mathbb{R} \right\}$$

- a) Show that R is a ring and that I is an ideal in R .
- b) Show that R/I and $\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ are isomorphic rings.
- c) Is R a semisimple ring? Is R/I a semisimple ring?
- d) Find 3 minimal left ideals in R .

Problem 2

Let R be a ring and M an R -module. Let N and L be submodules of M .

- a) Show that $N \cap L$ is a submodule of M , and give an example to show that $N \cup L$ is not always a submodule of M .
- b) Assume that M is a noetherian R -module. Show that then N and M/N are noetherian R -modules.

Problem 3

- a) Find the possible invariant factors and rational canonical forms for 6×6 -matrices over the real numbers \mathbb{R} , with minimal polynomial $(x^2 + 1)(x - 3)^2$.
- b) Let V be a vector space of dimension 4 over the real numbers \mathbb{R} , and let $T : V \rightarrow V$ be a linear transformation. Let v_1, v_2 be elements in V such that $\{v_1, v_2, Tv_2, T^2v_2\}$ is a basis for V , and assume that $Tv_1 = 2v_1$ and $T^3v_2 = 2v_2 + 3Tv_2$. In the usual way we view V (together with $T : V \rightarrow V$) as an $\mathbb{R}[x]$ -module. Show that v_1 and v_2 generate V as an $\mathbb{R}[x]$ -module, and find $f_1(x)$ and $f_2(x)$ in $\mathbb{R}[x]$, where $f_1(x) | f_2(x)$, such that $V \simeq \mathbb{R}[x]/(f_1(x)) \oplus \mathbb{R}[x]/(f_2(x))$.



Faglig kontakt under eksamen:
Petter Andreas Berg (73 59 04 83)

EXAM IN RINGS AND MODULES (MA3201)

Thursday, 9th December 2004

Time: 09:00 – 13:00

Grades to be announced: Thursday, 6th January 2005

Permitted aids: None.

Problem 1 Let

$$R = \left\{ \begin{pmatrix} a & 0 & 0 \\ b & a & 0 \\ c & d & e \end{pmatrix} \mid a, b, c, d, e \in \mathbb{C} \right\}.$$

a) Show that R is a ring under the usual addition and multiplication of matrices.

b) Let

$$I = \left\{ \begin{pmatrix} 0 & 0 & 0 \\ b & 0 & 0 \\ c & d & 0 \end{pmatrix} \mid b, c, d \in \mathbb{C} \right\}.$$

Show that I is a two-sided ideal in R , and that I is nilpotent.

c) Show that R/I and $\mathbb{C} \oplus \mathbb{C}$ are isomorphic rings. Is R/I a semisimple ring?

d) How can the two-sided ideals in the ring R/I be described in terms of two-sided ideals in R ? Find two maximal two-sided ideals in R .

Problem 2

a) Let $\varphi: R \rightarrow S$ be a homomorphism of rings. Show that any left S -module M becomes a left R -module by defining

$$r \cdot m = \varphi(r)m$$

for all r in R and m in M .

Recall the following: Let F be a field. Suppose A is an algebra over F ; that is, there is a map $F \times A \rightarrow A$, written $(\alpha, r) \mapsto \alpha \cdot r$, such that A is a vector space over F and

$$\alpha \cdot (rr') = (\alpha \cdot r)r' = r(\alpha \cdot r')$$

for all α in F , and all r and r' in A .

Assume that $0 \neq 1_A$ in A , where 1_A is the identity in A .

- b) Show that $\psi: F \rightarrow A$ given by $\psi(\alpha) = \alpha \cdot 1_A$, is a homomorphism of rings with $\text{Im } \psi \subseteq Z(A)$. Here

$$Z(A) = \{z \in A \mid za = az \text{ for all } a \in A\}.$$

Also, show that ψ is injective.

- c) Suppose that A is a finite dimensional algebra over F ; that is, $\dim_F A$ is finite. Show that A is both left artinian and left noetherian.

Let M be a finitely generated left A -module. Show that M is both an artinian and a noetherian A -module.

Problem 3 Let V be a vector space over a field F with $\dim_F V = n < \infty$. Let $T: V \rightarrow V$ be a non-zero linear transformation. Then V becomes an $F[x]$ -module by letting

$$x^i \cdot v = T^i(v)$$

for all v in V and $i \geq 0$. It is not necessary to prove this.

- a) Let $\text{Ann}_{F[x]} V = \{g(x) \in F[x] \mid g(x) \cdot v = 0 \text{ for all } v \in V\}$. Show that $\text{Ann}_{F[x]} V$ is an ideal in $F[x]$.

Let $f(x)$ be the minimal polynomial of T . Show that $\text{Ann}_{F[x]} V = (f(x))$.

- b) Suppose that T is a non-zero nilpotent linear transformation; that is, $T^l = 0$ for some positive integer l . Show that the minimal polynomial $f(x)$ of T is equal to x^m for some integer m with $0 < m \leq n$.
- c) Suppose also here that T is a non-zero nilpotent linear transformation. What is the smallest possible dimension of the kernel of T ? And what is the largest possible dimension of the kernel of T ?



Contact during exam: Øyvind Solberg/Petter Andreas Bergh
Telephone: 73 59 17 48/73 59 04 83

Exam in course MA3201 Rings and modules

English

Wednesday November 30, 2005

Time: 09.00-13.00

Permitted aids: none

Grades: 21.12.2005.

Problem 1 Let q be a fixed non-zero element in \mathbb{C} , the set of complex numbers. Define the subset R_q of the ring of 4×4 -matrices over \mathbb{C} by

$$R_q = \left\{ \begin{pmatrix} a & 0 & 0 & 0 \\ b & a & 0 & 0 \\ c & 0 & a & 0 \\ d & c & -qb & a \end{pmatrix} \mid a, b, c, d \in \mathbb{C} \right\}.$$

- a) Show that R_q is a ring.
- b) For which q in \mathbb{C} is R_q a commutative ring?
- c) For a given element α in \mathbb{C} define the subset

$$I_\alpha = \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 \\ b & 0 & 0 & 0 \\ \alpha b & 0 & 0 & 0 \\ d & \alpha b & -qb & 0 \end{pmatrix} \mid b, d \in \mathbb{C} \right\}$$

of R_q . Show that I_α is a left ideal in R_q for all α in \mathbb{C} .

- d) Show that each of the left ideals I_α is generated by one element as a left ideal. Show that $I_\alpha \simeq R/I_{\alpha q}$ as left R -modules.

Problem 2 Let \mathbb{Q} be the field of rational numbers, and let a and b in \mathbb{Q} be different elements. Find all possible rational canonical forms for 4×4 -matrices over \mathbb{Q} having

$$(x + a)^2(x + b)$$

as a minimal polynomial.

Problem 3 Let \mathbb{C} be the field of complex numbers and $\mathbb{C}[x]$ the polynomial ring over \mathbb{C} in one variable x . Let $\alpha \in \mathbb{C}$ be a complex number.

a) Show that the map $\varphi_\alpha: \mathbb{C}[x] \rightarrow \mathbb{C}$ defined by $\varphi_\alpha(f(x)) = f(\alpha)$ is a surjective ring homomorphism, and use this to show that the ideal generated by $x - \alpha$ is a maximal ideal in $\mathbb{C}[x]$.

b) For which $n \geq 1$ is the ring

$$\begin{pmatrix} \frac{\mathbb{C}[x]}{((x-\alpha)^n)} & \frac{\mathbb{C}[x]}{((x-\alpha)^n)} \\ \frac{\mathbb{C}[x]}{((x-\alpha)^n)} & \frac{\mathbb{C}[x]}{((x-\alpha)^n)} \end{pmatrix}$$

semisimple?

Problem 4 Let R be a ring, and let M be a Noetherian left R -module. Show that any surjective R -homomorphism $f: M \rightarrow M$ is an isomorphism. (Hint: Consider the chain $\text{Ker } f \subseteq \text{Ker}(f^2) \subseteq \text{Ker}(f^3) \subseteq \dots$ of submodules of M).



Contact during exam: Øyvind Solberg
Telephone: 73 59 17 48

EXAM IN RINGS AND MODULES (MA3201)

English
Friday 15th December 2006
Time: 09:00–13:00
Permitted aids: None

Grades: 15.01.2007.

Problem 1 Let A be the 3×3 matrix

$$\begin{pmatrix} 1 & 2 & -4 \\ 1 & 2 & 2 \\ -1 & 1 & 1 \end{pmatrix}$$

over \mathbb{C} , the complex numbers.

- a) Find the Smith normal form of the matrix $A - xI_3$ over the ring $\mathbb{C}[x]$, where $\mathbb{C}[x]$ is the polynomial ring in one variable x over \mathbb{C} and I_3 is the 3×3 identity matrix.
- b) Find the rational canonical form of the matrix A over \mathbb{C} .
- c) Find the Jordan canonical form of the matrix A over \mathbb{C} .

Problem 2 Let R and S be two rings. An abelian group M is called a S - R -bimodule if M is a left S -module and a right R -module, such that

$$s(mr) = (sm)r$$

for all s in S , for all r in R and for all m in M . Let

$$\Lambda = \begin{pmatrix} R & 0 \\ M & S \end{pmatrix}$$

where M is a S - R -bimodule different from (0) . Let $\begin{pmatrix} r & 0 \\ m & s \end{pmatrix}$ and $\begin{pmatrix} r' & 0 \\ m' & s' \end{pmatrix}$ be two elements in Λ . The set Λ becomes an abelian group under the binary operation, $+$, given by

$$\begin{pmatrix} r & 0 \\ m & s \end{pmatrix} + \begin{pmatrix} r' & 0 \\ m' & s' \end{pmatrix} = \begin{pmatrix} r+r' & 0 \\ m+m' & s+s' \end{pmatrix}.$$

Define a binary operation, \cdot , on Λ by letting

$$\begin{pmatrix} r & 0 \\ m & s \end{pmatrix} \cdot \begin{pmatrix} r' & 0 \\ m' & s' \end{pmatrix} = \begin{pmatrix} rr' & 0 \\ mr' + sm' & ss' \end{pmatrix}.$$

- a) Show that Λ is a ring with 1, when addition, $+$, and multiplication, \cdot , is defined as above.
- b) Find
 - (i) an idempotent element different from 0 and 1 in Λ ,
 - (ii) a nilpotent element different from 0 in Λ .
- c) Let $I = \left\{ \begin{pmatrix} 0 & 0 \\ m & 0 \end{pmatrix} \mid m \in M \right\}$. Show that I is a two-sided ideal in Λ . Show that $\Lambda/I \simeq R \oplus S$ as rings.

Problem 3 Let k be a field. The map $\varphi: k[x]/(x^2) \rightarrow k$ given by

$$\varphi(f(x) + (x^2)) = f(0)$$

is a homomorphism of rings. Let $R = k$ and $S = k[x]/(x^2)$.

- a) Let M be a left R -module. Show that M becomes a left S -module by defining

$$s \cdot m = \varphi(s)m$$

for all s in S and for all m in M .

- b) Let $M = k^2 = \{(a, b) \mid a, b \in k\}$. Then is M a left k -module by letting

$$\alpha(a, b) = (\alpha a, \alpha b)$$

and a right k -module by letting

$$(a, b)\alpha = (a\alpha, b\alpha)$$

for all α in k and for all (a, b) in M . With these module structures M becomes a k - k -bimodule (Do not need to show this). By a) we have that the left k -module M is a left S -module by letting $(f(x) + (x^2)) \cdot m = \varphi(f(x) + (x^2))m$. Show that M is a S - R -bimodule, when $R = k$ and $S = k[x]/(x^2)$.

- c) Now let $\Lambda = \begin{pmatrix} R & 0 \\ M & S \end{pmatrix}$, where M is as in b), and Λ is a ring as given in Problem 2. Show that Λ is an algebra over k . What is $\dim_k \Lambda$? Decide if Λ is
- (i) a left artinian ring,
 - (ii) a left noetherian ring,
 - (iii) a semisimple ring.
- d) Let J be the left ideal $\left\{ \begin{pmatrix} 0 & 0 \\ (0,a) & bx+(x^2) \end{pmatrix} \mid a, b \in k \right\}$. Consider the left Λ -module $X = \Lambda/J$. Show that $f: X \rightarrow X$ given by

$$f(\lambda + J) = \lambda \begin{pmatrix} 0 & 0 \\ (0,0) & 1+(x^2) \end{pmatrix} + J$$

is a Λ -homomorphism. Find the image $\text{Im } f$ of f . Show that $X = \text{Im } f \oplus Y$ for a submodule Y of X .



Faglig kontakt under eksamen:
Idun Reiten (99 24 45 39)

EXAM IN RINGS AND MODULES (MA3201)

Tuesday, 11.th December 2007

Time: 09:00 – 13:00

Grades to be announced: Friday, 21 December 2007

Permitted aids: None.

You should give a reason for all answers.

Problem 1

Let F be a field, R the matrix ring

$$R = \begin{pmatrix} F & 0 & 0 \\ F & F & 0 \\ F & F & F \end{pmatrix}$$

and

$$I = \begin{pmatrix} 0 & 0 & 0 \\ F & 0 & 0 \\ F & F & 0 \end{pmatrix}$$

- Show that I is an ideal in R , and that I is nilpotent. Is R a semisimple ring?
- Show that the factor ring R/I is a semisimple ring.
- Find 2 different minimal left ideals in R which are isomorphic as R -modules.

- d) Find all the ideals in R which contain the ideal I .

Problem 2

Let \mathbb{Z} be the ring of integers, and let R be the ring $\begin{pmatrix} \mathbb{Z} & 0 \\ \mathbb{Z} & \mathbb{Z} \end{pmatrix}$

- a) Find all idempotent elements in R , and describe the left ideals of the form Re for an idempotent element e in R .
- b) Let I be the left ideal $\begin{pmatrix} \mathbb{Z} & 0 \\ \mathbb{Z} & 0 \end{pmatrix}$. Find an infinite number of left ideals J in R such that $R = I \oplus J$.

Problem 3

- a) Show that the ring of integers \mathbb{Z} is noetherian, and not artinian.
- b) Give a proof of the fact that if M is a noetherian module over a ring R , then M is a finitely generated R -module.

Problem 4

- a) Denote by \mathbb{R} the real numbers. Find the Smith normal form over $\mathbb{R}[x]$ for the matrix

$$\begin{pmatrix} -3-x & 2 & 0 \\ 1 & -x & 1 \\ 1 & -3 & -2-x \end{pmatrix}$$

Let $V = \mathbb{R}^3$, and let $T = T_A : V \rightarrow V$ be the linear transformation given by the matrix $A = \begin{pmatrix} -3 & 2 & 0 \\ 1 & 0 & 1 \\ 1 & -3 & -2 \end{pmatrix}$ with respect to the standard basis for $V = \mathbb{R}^3$. Describe the $\mathbb{R}[x]$ -module V (defined using $T : V \rightarrow V$) in terms of cyclic $\mathbb{R}[x]$ -modules.

- b) Let A be a 7×7 matrix over \mathbb{R} , with characteristic polynomial $c(x) = -(x-1)^2(x-2)^3(x^2+1)$ and with minimal polynomial $m(x)$ of degree 5. Find all the possibilities for the invariant factors for A , (that is, for $xI - A$), and in each case, the associated rational canonical form for A .