TMA4100 Øving 1

August 31, 2011

Exercise 1.5.59

Find a formula for the inverse function f^{-1} and verify that $(f \circ f^{-1})(x) = (f^{-1} \circ f)(x) = x$.

(a)

$$f(x) = \frac{100}{1 + 2^{-x}}$$

(b)

$$f(x) = \frac{50}{1+1.1^{-x}}$$

Exercise 2.1.3

Find the average rate of change of the function $h(t) = \cot t$ over the intervals

(a)
$$[\pi/4, 3\pi/4]$$
, and

(b)

 $[\pi/6,\pi/2]$.

Exercise 2.3.20

We have

$$f(x) = \sqrt{x-7}, \quad L = 4, \quad x_0 = 23, \quad \epsilon = 1.$$

First, find an open interval about x_0 on which the inequality $|f(x) - L| < \epsilon$ holds. Then give a value for $\delta > 0$ such that for all x satisfying $0 < |x - x_0| < \delta$ the inequality $|f(x) - L| < \epsilon$ holds.

Exercise 2.3.53

Show by example that the following statement is wrong: The number L is the limit of f(x) as x approaches x_0 if f(x) gets closer to L as x approaches x_0 .

Explain why the function in your example does not have the given value of L as a limit as $x \to x_0$.

Exercise 2.5.35

Graph the rational function

$$y = \frac{x^2 - 4}{x - 1}$$
.

Include the graphs and equations of its asymptotes.

Exercise 2.6.39

For what value of a is

$$f(x) = \begin{cases} x^2 - 1, & x < 3\\ 2ax, & x \ge 3 \end{cases}$$

continuous at every x?

Exercise 2.6.46

Explain why the equation $\cos x = x$ has at least one solution.

Exercise 2.7.34

Does the graph

$$g(x) = \begin{cases} x \sin(1/x), & x \neq 0\\ 0, & x = 0 \end{cases}$$

have a tangent at the origin? Give reasons for your answer.

Exercise 3.2.53

The curve $y = ax^2 + bx + c$ passes through the point (1, 2) and is tangent to the line y = x at the origin. Find a, b and c.

Exercise 3.3.13

Had Galileo dropped a cannonball from the Tower of Pisa, 179 ft above the ground, the ball's height above the ground t sec into the fall would have been

$$s = 179 - 16t^2$$
.

- a) What would have been the ball's velocity, speed and acceleration at time t?
- b) About how long would it have taken the ball to hit the ground?
- c) What would have been the ball's velocity at the moment of impact?

Exercise 3.4.25

Find y'' if

1. $y = \csc x$.

2. $y = \sec x$.

Exercise 3.5.97

Find a parametrization for the lower half of the parabola $x - 1 = y^2$.

Exercise 3.6.2

Use implicit differentiation to find dy/dx when

$$x^3 + y^3 = 18xy \; .$$

Exercise 3.6.32

Verify that the point (-2, 1) is on the curve

$$y^2 - 2x - 4y - 1 = 0 ,$$

and find the lines that are (a) tangent and (b) normal to the curve at the given point.

Exercise 3.7.9

Suppose that the differentiable function y = f(x) has an inverse and that the graph of f passes through the point (2, 4) and has a slope of 1/3 there. Find the value of df^{-1}/dx at x = 4.

Exercise 3.8.13

Find the limit

$$\lim_{x \to 1^-} \sin^{-1} x \; .$$