

# TMA 4115 Matematikk 3

## Lecture for KJ & NANO

Alexander Schmeding

NTNU

23. March 2017

In today's lecture we will...

- Recap: Similarity of matrices
- use similarity to relate matrices to diagonal matrices
- complex eigenvalues and eigenvectors

# Similarity of matrices

## Similarity of matrices $A$ and $B$

$A$  is **similar** to  $B$  if there is an invertible matrix  $P$  with

$$PAP^{-1} = B \quad (\text{or equivalently } A = P^{-1}BP)$$

Recall that similar matrices have the same characteristic polynomial (hence the same eigenvalues).

- $A = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$  and  $B = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$  are not similar.
- $A = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$  and  $C = \begin{bmatrix} 0 & 1 \\ -4 & 4 \end{bmatrix}$  are similar, since

$$C = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix}$$

We compute

- eigenvalues via the **characteristic polynomial**  $\det(A - \lambda I)$ .
- eigenvectors via  $(A - \lambda I)\vec{x} = \vec{0}$  (for  $\lambda$  eigenvalue!)

Example: Find eigenvalues and eigenvectors of  $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

Calculate the roots of the **characteristic polynomial**

$$\det(A - \lambda I) = (1 - \lambda)^2 - 1 = \lambda^2 - 2\lambda = (0 - \lambda)(2 - \lambda).$$


Roots of the characteristic polynomial = eigenvalues.

$\lambda = 0$  and  $\lambda = 2$ .

Calculate eigenvectors for  $\lambda = 0$  and  $\lambda = 2$  via Gaussian elimination, i.e. find a non trivial solution of  $(A - 0I)\vec{x} = \vec{0}$  and  $(A - 2I)\vec{y} = \vec{0}$ .

For example  $\vec{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  and  $\vec{y} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

Along the eigenvectors the matrix acts by scalar multiplication:



Observe that the eigenvectors even form a basis  $\mathcal{B}$  for  $\mathbb{R}^2$ !

## Observation: Matrix action in $\mathcal{B}$ coordinates

Recall  $\mathcal{B}$ -coordinates  $[\vec{x}]_{\mathcal{B}}$ , have invertible linear map ( $\sim$  matrix)  $P_{\mathcal{B}}$  with  $P_{\mathcal{B}}[\vec{x}]_{\mathcal{B}} = \vec{x}$ . Then

$$\begin{array}{ccc}
 \vec{x} & \xrightarrow{A} & A\vec{x} \\
 \downarrow P_{\mathcal{B}}^{-1} & & \uparrow P_{\mathcal{B}} \\
 [\vec{x}]_{\mathcal{B}} & \xrightarrow{\begin{bmatrix} 0 & 0 \\ 0 & 2 \end{bmatrix}} & [A\vec{x}]_{\mathcal{B}}
 \end{array}$$

$\rightarrow A$  acts on the  $\mathcal{B}$ -coordinates as a diagonal matrix!

Note  $A = P_{\mathcal{B}} \begin{bmatrix} 0 & 0 \\ 0 & 2 \end{bmatrix} P_{\mathcal{B}}^{-1}$

**Question:**

Can we “transform” (all?) square matrices to diagonal matrices?

A a  $n \times n$ -matrix. Find a similar diagonal matrix (if possible).

1. Compute the eigenvalues of  $A$ ,
2. Compute the eigenvectors associated to the eigenvalues.  
(Gaussian elimination!)
3. Check if we have  $n$  linear independent eigenvectors. If not:  $A$  is not diagonalisable!  
**Hint:** Eigenvectors of different eigenvalues are linearly independent.
4. Write eigenvectors as columns in a matrix  $P$
5. Construct a diagonal matrix  $D$  whose diagonal entries are the eigenvalues corresponding to the columns in  $P$
6. Then  $A = PDP^{-1}$