

# TMA 4115 Matematikk 3

## Lecture 4 for KJ & NANO

Alexander Schmeding

NTNU

19. January 2016

Topics in todays lecture...

- Recap: The complex exponential function
- 2nd order differential equations

# Complex functions

A *complex function*  $f: U \rightarrow V$ ,  $x \mapsto f(x)$  consists of  $U \subseteq \mathbb{C}$  (domain),  $V \subseteq \mathbb{C}$  (codomain) and a rule  $x \mapsto f(x)$ .

## Complex polynomials

$p: \mathbb{C} \rightarrow \mathbb{C}$ ,  $p(z) = \sum_{k=0}^n a_k z^k$  (complex) *polynomial*, where  $a_0, \dots, a_n \in \mathbb{C}$  and  $n \in \mathbb{N}$ .

$w \in \mathbb{C}$  is a root of  $p$  if  $p(w) = 0$ . For  $p(z) = az^2 + bz + c$  we can find roots by the quadratic formula.

# The complex exponential function

$\exp: \mathbb{C} \rightarrow \mathbb{C}$ ,  $\exp(x + iy) = e^x(\cos(y) + i \sin(y))$  write  
 $e^z := \exp(z)$

## Properties of $\exp$ for $z = x + iy$

- (a)  $\exp(a + 0i) = e^a$ , ( $\exp$  for real numbers),
- (b)  $|e^z| = e^x$ ,  $\arg(e^z) = y$ ,
- (c)  $\operatorname{Re}(e^z) = e^x \cos(y)$  and  $\operatorname{Im}(e^z) = e^x \sin(y)$ ,
- (d)  $e^{z_1+z_2} = e^{z_1} e^{z_2}$  for  $z_1, z_2 \in \mathbb{C}$ ,
- (e)  $\frac{1}{e^z} = e^{-z}$   $\overline{\exp(z)} = \exp(\bar{z})$ ,

Can use  $\exp$  to write polar coordinates:

$$z = r(\cos(\theta) + i \sin(\theta)) = re^{i\theta}$$

We know how to solve the differential equations

$$y'(t) = \frac{d}{dt}y(t) = f(y, t)$$

if the function  $f$  is “nice”.

This is a *first order differential equation*, because it **only involves the first derivative** of  $y$ .

First order differential equations are “simple”. The (physical) world is complicated, whence it can not only be described by first order equations. We need more complicated equations!

## 3.1 Newtons second law

### Newton's second law

The *acceleration*  $a$  of a body with *mass*  $m$  is proportional to the *net force*  $F$  via

$$F = ma \quad (1)$$

**Question:** What is the displacement  $y(t)$  of the body from a reference point?

Acceleration  $a$  is rate of change of velocity  $v$ , i.e.

$$a = \frac{d}{dt}v = v' \quad (2)$$

Velocity  $v$  is rate of change of the displacement, i.e.

$$v = \frac{d}{dt}y = y' \quad (3)$$

Thus (2) becomes  $a = y''$ .

What is the displacement  $y(t)$  of the body from a reference point?

Finally,  $F$  depends on time, displacement and velocity, thus

$$F(t, y, v) = F(t, y, y') \quad (\text{using (3)})$$

With Newtons second law (1) obtain

$$F(t, y, y') = my''.$$

This is a **second order differential equation** since it involves derivatives of  $y$  of up to second order.

## 3.2 Definition

A *second order differential equation* is an equation of the form

$$\frac{d^2}{dt^2}y(t) = f\left(y, \frac{d}{dt}y, t\right) \quad (4)$$

where  $f$  is a given function.


A *solution* to (4) is a function  $y$  which is twice continuously differentiable and satisfies (4).

Usually we write  $y' = \frac{d}{dt}y(t)$  and  $y'' = \frac{d^2}{dt^2}y(t)$ , thus (4) reads:

$$y'' = f(t, y, y') \quad (5)$$

### 3.3 The vibrating spring

We consider a spring suspended from a beam:



Attach weight  $m$ :  
 New equilibrium at height  $x_0$   
 Stretch and release the spring  
 ... it will move!

Forces acting on weight in motion:

**damping force**  $D(v)$  (depends on velocity  $v = x'$ ),  
**external force**  $F(t)$ , **Restoring force**  $R(x)$  and **gravity**  $mg$ .



## A model for the position $x(t)$ of the spring

### Newton's second law (1) for the spring

$$\begin{aligned} ma &= \text{total force acting on the weight} \\ &= R(x) + mg + D(v) + F(t) \end{aligned} \quad (6)$$

Velocity  $v = x'$  and acceleration  $a = v' = x''$ , thus (6) becomes

$$mx'' = R(x) + mg + D(x') + F(t). \quad (7)$$

### Hooke's law (valid for some springs)

$R(x) = -kx$  for  $k > 0$  constant and small  $x$ .

Assuming Hooke's law, (7) becomes

$$x'' = -\frac{k}{m}x + g + \frac{D(x') + F(t)}{m}. \quad (8)$$

# Is there a solution for every 2nd order differential equation?

## 3.4 Theorem

Let  $p, q$  and  $g$  be continuous functions on the interval  $(\alpha, \beta)$ . Fix  $t_0 \in (\alpha, \beta)$  and  $y_0, y_1 \in \mathbb{R}$ . There is a unique function  $y: (\alpha, \beta) \rightarrow \mathbb{R}$  which solves

$$\begin{cases} y'' + p(t)y' + q(t)y = g(t), & \text{for } t \in (\alpha, \beta) \\ y(t_0) = y_0, y'(t_0) = y_1 \end{cases} \quad (9)$$

## 3.5 Remark

The solution of (9) is defined on all of  $(\alpha, \beta)$

We need  $y(t_0) = y_0, y'(t_0) = y_1$  to get a unique solution.

**Open Problem:** How to find a solution?