

Summary Lecture 24: Residue integration

- ① **Residues:** $\operatorname{Res}_{z=z_0} f(z) = b_1$ where

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n + \frac{b_1}{z - z_0} + \frac{b_2}{(z - z_0)^2} + \dots, \quad 0 < |z - z_0| < R$$

z_0 order 1 pole:

$$\operatorname{Res}_{z=z_0} f(z) = \lim_{z \rightarrow z_0} (z - z_0) f(z)$$

$$\operatorname{Res}_{z=z_0} \frac{p(z)}{q(z)} = \frac{p(z_0)}{q'(z_0)}$$

z_0 order n pole:

$$\operatorname{Res}_{z=z_0} f(z) = \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \left((z - z_0)^n f(z) \right)^{(n-1)}$$

z_0 isolated essential singularity: Find b_1 from the Laurent series!

- ② **Residue theorem:** Assume

(A1) C simple, closed curve oriented counterclockwisely

(A2) $f(z)$ has finite number of singularities z_1, \dots, z_m enclosed by C

Then
$$\oint_C f(z) dz = 2\pi i \sum_{j=1}^m \operatorname{Res}_{z=z_j} f(z)$$

Lecture 25: Residue integration of real integrals

Kreyszig: Section 16.4

- 1 Computing real integrals:

Type I: $\int_0^{2\pi} F(\cos \theta, \sin \theta) d\theta$

Type II: $\int_{-\infty}^{\infty} f(x) dx$

Type III: $\int_{-\infty}^{\infty} f(x) e^{-iwz} dx$

- 2 Examples

Information (check [www](#)):

Øving 13, 8/13 øvinger for å ta eksamen, neste uke

Frist øving 12 og 13:

Denne uka! Onsdag kl 23:59!

Lecture 25: Type I real integrals

$$I = \int_0^{2\pi} F(\cos \theta, \sin \theta) d\theta, \quad F \text{ rational}$$

Substitution:

$$z = e^{i\theta}$$

$$\cos \theta = \frac{1}{2}\left(z + \frac{1}{z}\right), \quad \sin \theta = \frac{1}{2i}\left(z - \frac{1}{z}\right)$$

$$dz = iz d\theta$$


Segment $[0, 2\pi)$ \rightsquigarrow circle $|z| = 1$

Lecture 25: Type II real integrals

$$I = \int_{-\infty}^{\infty} f(x) dx, \quad f \text{ rational, no real singularities}$$

$$(i) I = \lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx$$

$$(ii) \int_{-R}^R f(x) dx = \oint_{C_R} f(z) dz - \int_{S_R} f(z) dz$$



(iii) R big enough $\Rightarrow C_R$ encircles all poles in upper half plane

$$\oint_{C_R} f(z) dz \stackrel{\text{Residue thm.}}{=} 2\pi i \sum_{\substack{\text{poles in upper} \\ \text{half plane}}} \text{Res } f(z)$$

$$(iv) \left| \int_{S_R} f(z) dz \right| \leq \max_{z \in S_R} |f(z)| \cdot L \leq \max_{|z|=R} |f(z)| \cdot \pi R \xrightarrow[R \rightarrow \infty]{\text{Must show}} 0$$


Lecture 25: Type III real integrals

$$I = \int_{-\infty}^{\infty} f(x)e^{-iwx} dx, \quad f \text{ rational, no real sing.}$$


Same procedure as for Type II integrals. . .

. . . and choose C_R to insure $w \cdot \text{Im } z \leq 0$:

$$w \leq 0 \rightsquigarrow$$



$$; \quad w > 0 \rightsquigarrow$$



$$\implies |f(z)e^{-i wz}| \stackrel{z=x+iy}{=} |f(z)|e^{w \cdot y} \stackrel{z \in C_R}{\leq} |f(z)|$$

$$\text{Re } I = \int_{-\infty}^{\infty} f(x) \cos(wx) dx$$

$$\text{Im } I = - \int_{-\infty}^{\infty} f(x) \sin(wx) dx$$

Summary Lecture 25: Real integrals


1 Type I: $I = \int_0^{2\pi} F(\cos \theta, \sin \theta) d\theta$ F rational

Substitution: $z = e^{i\theta}$, $\cos \theta = \frac{1}{2}(z + \frac{1}{z}), \dots$, $[0, 2\pi) \rightsquigarrow |z| = 1$

2 Type II: $I = \int_{-\infty}^{\infty} f(x) dx$ f rational, no real singularities, ...

(i) $I = \lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx$

(ii) $\int_{-R}^R f(x) dx = \oint_{C_R} f(z) dz - \int_{S_R} f(z) dz$



(iii) R big enough $\Rightarrow C_R$ encircles all poles in upper half plane

$$\oint_{C_R} f(z) dz \stackrel{\text{Residue thm.}}{=} 2\pi i \sum_{\text{poles in upper half plane}} \text{Res } f(z)$$


(iv) $|\int_{S_R} f(z) dz| \leq \max_{z \in S_R} |f(z)| \cdot L \leq \max_{|z|=R} |f(z)| \cdot \pi R \xrightarrow{R \rightarrow \infty} 0$ Must show 0

Summary Lecture 25: Real integrals

- ③ **Type III:** $I = \int_{-\infty}^{\infty} f(x)e^{-iwx} dx$ f rational, no real singularities, ...

Same procedure as for Type II integrals. . .

...and choose C_R to insure $w \cdot \text{Im } z \leq 0$:



$$\implies |f(z)e^{-iwz}| \stackrel{z=x+iy}{=} |f(z)|e^{w \cdot y} \stackrel{z \in C_R}{\leq} |f(z)|$$

- ④ $\text{Re } I = \int_{-\infty}^{\infty} f(x) \cos(wx) dx$ $\text{Im } I = - \int_{-\infty}^{\infty} f(x) \sin(wx) dx$