

# Laplace Transform

①  $F(s) = \mathcal{L}[f](s) = \int_0^\infty f(t)e^{-st}dt$

## ② Properties:

Linearity:  $\mathcal{L}[af(t) + bg(t)](s) = a\mathcal{L}[f](s) + b\mathcal{L}[g](s)$

s-shift:  $\mathcal{L}[e^{at}f(t)](s) = \mathcal{L}[f](s - a)$  for  $s - a > k$

t-shift:  $\mathcal{L}[f(t - a)u(t - a)](s) = e^{-as}\mathcal{L}[f](s)$

Derivatives:  $\mathcal{L}[f'(t)](s) = s\mathcal{L}[f](s) - f(0)$

Integral:  $\mathcal{L}[\int_0^t f(\tau)d\tau](s) = \frac{1}{s}\mathcal{L}[f](s)$

Convolution:  $\boxed{\mathcal{L}[f * g](s) = \mathcal{L}[f](s)\mathcal{L}[g](s)}$   $(f * g)(t) = \int_0^t f(\tau)g(t - \tau)d\tau$

## ③ Unit step function, delta function:

$\delta$ -function:  $\int_0^\infty \delta(t - a)f(t)dt = f(a)$  for all  $f$  cont. at  $t = a$ .

$$\mathcal{L}[u(t - a)] = \frac{1}{s}e^{-as}, \quad \boxed{\mathcal{L}[\delta(t - a)] = e^{-as}}$$

# Lecture 4: Laplace Transform

Kreyszig: Sections 6.6, 6.7, 11.1

- ① Differentiation of transforms
- ② Integration of transforms
- ③ Systems of differential equations
- ④ Fourier series (introduction)
- ⑤ Examples

## Lecture 4: Laplace Transform


Derivatives and integrals of transforms:

$$-F'(s) = \mathcal{L}[tf(t)](s)$$

$$\int_s^\infty F(\bar{s})d\bar{s} = \mathcal{L}\left[\frac{1}{t}f(t)\right](s)$$

# Lecture 4: Laplace Transform

Systems of ordinary differential equations:



## Lecture 4: Fourier Analysis

$f(x)$  is  $p$ -periodic if  $f(x + p) = f(x)$  for all  $x \in \mathbb{R}$ .

Representation of  $f(x)$  by trigonometric series:

$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx).$$

# Summary: Laplace Transform

①  $F(s) = \mathcal{L}[f](s) = \int_0^\infty f(t)e^{-st}dt$

## ② Properties:

Linearity:  $\mathcal{L}[af(t) + bg(t)](s) = a\mathcal{L}[f](s) + b\mathcal{L}[g](s)$

s-shift:  $\mathcal{L}[e^{at}f(t)](s) = \mathcal{L}[f](s-a)$  for  $s-a > k$

t-shift:  $\mathcal{L}[f(t-a)u(t-a)](s) = e^{-as}\mathcal{L}[f](s)$

Derivatives:  $\mathcal{L}[f'(t)](s) = s\mathcal{L}[f](s) - f(0)$

$$\boxed{\mathcal{L}^{-1}[F'(s)](t) = -tf(t)}$$

Integral:  $\mathcal{L}[\int_0^t f(\tau)d\tau](s) = \frac{1}{s}\mathcal{L}[f](s)$

$$\boxed{\mathcal{L}^{-1}[\int_s^\infty F(\bar{s})d\bar{s}](t) = \frac{1}{t}f(t)}$$

Convolution:  $\mathcal{L}[f * g](s) = \mathcal{L}[f](s)\mathcal{L}[g](s)$   $(f * g)(t) = \int_0^t f(\tau)g(t-\tau)d\tau$

③ Unit step and delta function:  $\mathcal{L}[u(t-a)] = \frac{1}{s}e^{-as}$ ,  $\mathcal{L}[\delta(t-a)] = e^{-as}$