

Complex Analysis

1 Sets in \mathbb{C} :

Open: Contains open ball about each point

Connected: Any two points can be connected by a finite continuous curve within the set

Domain: Open and connected




2 Complex functions:

Assigns each z in the domain of definition a unique value $f(z) \in \mathbb{C}$

$$w = f(z) = u(x, y) + iv(x, y)$$

$$u = \operatorname{Re} w, \quad v = \operatorname{Im} w$$



3 **Limit, continuity:** Same as for functions of 2 real variables

4 **Derivative:** Same definition/rules as for functions of one real variable

5 **Analytic functions:**

$f(z)$ **analytic** in domain D if defined and *differentiable* in all $z \in D$

\Leftrightarrow **Cauchy-Riemann equations** hold in D :
$$u_x = v_y, \quad u_y = -v_x$$

Lecture 15: Complex Analysis

Kreyszig: Sections 13.4, 17.1

- 1 Cauchy-Riemann equations
- 2 Laplace equation, harmonic functions
- 3 Conformal mappings

OBS: Du trenger 8 (av 12) øvinger godkjent for å få ta eksamen!!

Lecture 15: The Cauchy-Riemann equations

$$(CR) \quad \boxed{u_x = v_y} \quad \text{and} \quad \boxed{u_y = -v_x}$$

$f(z) = u(x, y) + iv(x, y)$ analytic in domain D



u_x, u_y, v_x, v_y exists, continuous, and satisfy (CR) in D

Lecture 15: Laplace equation

(Laplace) $u_{xx} + u_{yy} = 0$

$f(z) = u(x, y) + iv(x, y)$ analytic in domain D



u, v are 2x cont. differentiable + satisfy (Laplace) in D

$\rightarrow u, v$ are conjugate harmonic functions

Lecture 15: Conformal mappings

Preserves angles and orientation between smooth curves

$$f \text{ analytic in } D \Rightarrow f \text{ conformal where } f' \neq 0 \text{ in } D$$

Ex: $f(z) = z^n$ conformal at $z \neq 0$

$$\text{At } z = 0: \arg(\dot{w}_1 - \dot{w}_2) = n \arg(\dot{z}_1 - \dot{z}_2)$$

Summary Lecture 15: Complex Analysis

1 Analytic functions and Cauchy-Riemann equations:

$$f(z) = u(x, y) + iv(x, y) \text{ analytic in domain } D$$



u_x, u_y, v_x, v_y exists, are continuous, and

$$\boxed{u_x = v_y \quad \text{and} \quad u_y = -v_x} \quad \text{in} \quad D.$$

2 Laplace equation $u_{xx} + u_{yy} = 0$

$$f(z) = u(x, y) + iv(x, y) \text{ analytic in domain } D$$



u, v are 2 times continuously differentiable, and

$$\boxed{u_{xx} + u_{yy} = 0 \quad \text{and} \quad v_{xx} + v_{yy} = 0} \quad \text{in} \quad D.$$

u, v are **conjugate harmonic** functions.

3 Conformal mappings:

Maps preserving angles and orientation between smooth curves

$$\boxed{f \text{ analytic in } D \Rightarrow f \text{ conformal where } f' \neq 0 \text{ in } D}$$

$$f(z) = z^n \text{ conformal at } z \neq 0, \quad z = 0: \arg(\dot{w}_1 - \dot{w}_2) = n \arg(\dot{z}_1 - \dot{z}_2)$$