

Formlar i numerikk

- La $p(x)$ vere eit polynom av grad $\leq n$ som interpolerer $f(x)$ i punkta $x_i, i = 0, 1, \dots, n$. Dersom x og alle nodane ligg i intervallet $[a, b]$, har vi at

$$f(x) - p(x) = \frac{1}{(n+1)!} f^{(n+1)}(\xi) \prod_{i=0}^n (x - x_i)$$

Dersom nodane er jamnt fordelte (inkludert endepunktane), og $|f^{(n+1)}(x)| \leq M$, har vi at

$$|f(x) - p(x)| \leq \frac{1}{4(n+1)} M \left(\frac{b-a}{n} \right)^{n+1}$$

- Numerisk derivasjon:

$$f'(x) = \frac{1}{h}(f(x+h) - f(x)) - \frac{h}{2}f''(\xi)$$

$$f'(x) = \frac{1}{h}(f(x) - f(x-h)) + \frac{h}{2}f''(\xi)$$

$$f'(x) = \frac{1}{2h}(f(x+h) - f(x-h)) + \frac{h^2}{2}f'''(\xi)$$

$$f''(x) = \frac{1}{h^2}(f(x+h) - 2f(x) + f(x-h)) - \frac{1}{12}h^2f^{(4)}(\xi)$$

- Newtons metode for likningssystemet $\mathbf{f}(\mathbf{x}) = \mathbf{0}$ er gitt ved

$$\begin{aligned} \mathbf{J}^{(k)} \cdot \Delta \mathbf{x}^{(k)} &= -\mathbf{f}(\mathbf{x}^{(k)}) \\ \mathbf{x}^{(k+1)} &= \mathbf{x}^{(k)} + \Delta \mathbf{x}^{(k)} \end{aligned}$$

- Iterative teknikkar for løysing av eit lineært likningssystem

$$\sum_{j=1}^n a_{ij}x_j = b_i, \quad i = 1, 2, \dots, n$$

$$\text{Jacobi : } x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij}x_j^{(k)} - \sum_{j=i+1}^n a_{ij}x_j^{(k)} \right)$$

$$\text{Gauss-Seidel : } x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij}x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij}x_j^{(k)} \right)$$

- Heuns metode for løysing av $\mathbf{y}' = \mathbf{f}(x, \mathbf{y})$:

$$\mathbf{K}_1 = h \mathbf{f}(x_n, \mathbf{y}_n)$$

$$\mathbf{K}_2 = h \mathbf{f}(x_n + h, \mathbf{y}_n + \mathbf{K}_1)$$

$$\mathbf{y}_{n+1} = \mathbf{y}_n + \frac{1}{2}(\mathbf{K}_1 + \mathbf{K}_2)$$

Sjå og formlane i Rottmann.

Tabell over Laplacetransformerte

$f(t)$	$\mathcal{L}(f)$
1	$\frac{1}{s}$
t	$\frac{1}{s^2}$
t^n ($n = 0, 1, 2, \dots$)	$\frac{n!}{s^{n+1}}$
e^{at}	$\frac{1}{s - a}$
$\cos \omega t$	$\frac{s}{s^2 + \omega^2}$
$\sin \omega t$	$\frac{\omega}{s^2 + \omega^2}$
$\cosh at$	$\frac{s}{s^2 - a^2}$
$\sinh at$	$\frac{a}{s^2 - a^2}$
$e^{at} \cos \omega t$	$\frac{s - a}{(s - a)^2 + \omega^2}$
$e^{at} \sin \omega t$	$\frac{\omega}{(s - a)^2 + \omega^2}$