

Løsningsforslag
Eksamens 06.08.99 i SIF5013/14

Oppgave 1

$$c_{n+2} = \frac{-n^2 + 5n - 6}{(n+2)(n+1)} = \frac{-(n-2)(n-3)}{(n+2)(n+1)}$$

$$c_0 = c_1 = 1, \quad c_2 = -3, \quad c_3 = -1/3$$

$$\underline{\underline{y = 1 + x - 3x^2 - \frac{1}{3}x^3}}$$

Oppgave 2

a)

$$\mathcal{L}^{-1} \left(\frac{as+b}{(x+2)^2+4} \right) = \mathcal{L}^{-1} \left(\frac{a(s+2)+b-2a}{(s+2)^2+4} \right)$$

$$= e^{-2t} \left(a \cos 2t + \left(\frac{b}{2} - a \right) \sin 2t \right)$$

$$\underline{\underline{\underline{y(t) = e^{-2t}(2\cos 2t + \frac{b}{2}\sin 2t) + \frac{1}{2}(t-2)^2u(t-2)}}}$$

b) Laplacetransformerer:

$$(s^2 + 4s + 8)Y - 2s - 9 = \frac{s^2 + 4s + 8}{s^3} e^{-2s}$$

$$Y(s) = \frac{2s+9}{(s+2)^2+4} + \frac{1}{s^3} e^{-2s}$$

$$\underline{\underline{y(t) = e^{-2t}(2\cos 2t + \frac{5}{2}\sin 2t) + \frac{1}{2}(t-2)^2u(t-2)}}$$

Oppgave 3

a) Setter inn:

$$h'(t) \sin ax = -\kappa h(t) a^2 \sin ax \Rightarrow h(t) = A e^{-\kappa a^2 t}$$

$$h(0) = 1 \Rightarrow \underline{\underline{h(t) = e^{-\kappa a^2 t}}}$$

b) Fourierrekken (sinusrekke p.g.a. odde funksjon)

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx = \frac{2}{\pi} \int_0^{\pi/2} 1 \cdot \sin nx dx = \frac{2}{\pi n} \left(1 - \cos \frac{\pi n}{2} \right)$$

Fra resultatet i a) følger

$$u(x, t) = \sum_{n=1}^{\infty} \frac{2}{\pi n} \left(1 - \cos \frac{\pi n}{2} \right) e^{-\kappa n^2 t} \sin nx$$

$$\underline{\underline{\underline{u(x, t) = \sum_{n=1}^{\infty} \frac{2}{\pi n} \left(1 - \cos \frac{\pi n}{2} \right) e^{-\kappa n^2 t} \sin nx}}}$$

Oppgave 4

a)

$$\begin{aligned}
 T(af_1 + bf_2) &= af_1(t) + bf_2(t) + \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{1}{1+(t-\tau)^2} (af_1(\tau) + bf_2(\tau)) d\tau \\
 &= a \left(f_1(t) + \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{1}{1+(t-\tau)^2} f_1(\tau) d\tau \right) \\
 &\quad + b \left(f_2(t) + \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{1}{1+(t-\tau)^2} f_2(\tau) d\tau \right) \\
 &= aTf_1 + bTf_2
 \end{aligned}$$

Fouriertransformerer:

$$\hat{g}(\omega) = \hat{f}(\omega) + \mathcal{F}\left(\frac{1}{\pi} \frac{1}{1+t^2}\right) \cdot \hat{f}(\omega) = \hat{f}(\omega) + e^{-|\omega|} \hat{f}(\omega) = (1 + e^{-|\omega|}) \hat{f}(\omega)$$

Transferfunksjon: $1 + e^{-|\omega|}$

(OBS: Feil i oppgaveteksten, $1 + e^{|\omega|}$, ble rettet på eksamen.)

b)

$$\begin{aligned}
 \hat{g}(\omega) &= (1 + e^{-|\omega|}) \hat{f}(\omega) \Rightarrow \hat{f}(\omega) = (i\omega) e^{-\omega^2/2} \\
 \underline{\underline{f(t)}} &= \frac{d}{dt} \left(\frac{1}{\sqrt{2\pi}} e^{-t^2/2} \right) = -\frac{1}{\sqrt{2\pi}} t e^{-t^2/2}
 \end{aligned}$$

Oppgave 5

a) Tabell over dividerte differanser:

x	f[]	f[,]	f[,,]
-1	-4		
0		3	
1	-1		-1
		1	
2	0	2	1
		5	
2	5		

Polynomet blir:

$$\begin{aligned}
 p(x) &= -4 + 3(x+1) - (x+1)x + (x+1)x + (x+1)x(x-1) \\
 &= \underline{\underline{x^3 - x^2 + x - 1}}
 \end{aligned}$$

b) La det nye polynomet være

$$q(x) = p(x) + c \cdot (x+1)(x-1)x(x-2)$$

Dette interpolerer datapunktene. I tillegg er

$$q^{(4)}(x) = 4!c = 24 \Rightarrow c = 1$$

og

$$\underline{\underline{q(x) = x^4 - x^3 - 2x^2 + 3x - 1}}$$

Oppgave 6

a) Sett $y_1 = x$, $y_2 = x'$

dvs. at systemet blir

$$y'_1 = y_2 \quad y_1(0) = 0$$

$$y'_2 = \cos(y_1) \quad y_2(0) = 1$$

b) Bruk av 3. ordens RK-metode gir:

1. skritt:

$$k_1 = \begin{bmatrix} 0.1 \\ 0.1 \end{bmatrix}, \quad k_2 = \begin{bmatrix} 0.11 \\ 0.09950 \end{bmatrix}, \quad y_1 = \begin{bmatrix} 0.1050 \\ 1.0998 \end{bmatrix} \approx y(0.1)$$

2. skritt:

$$k_1 = \begin{bmatrix} 0.10998 \\ 0.09945 \end{bmatrix}, \quad k_2 = \begin{bmatrix} 0.11992 \\ 0.09770 \end{bmatrix}, \quad y_2 = \begin{bmatrix} 0.2199 \\ 1.1983 \end{bmatrix} \approx y(0.2)$$

dvs:

$$\underline{\underline{x(0.2) \approx 0.2199.}}$$