

Formlar i numerikk

- La $p(x)$ vere eit polynom av grad $\leq n$ som interpolerer $f(x)$ i punkta $x_i, i = 0, 1, \dots, n$. Dersom x og alle nodane ligg i intervallet $[a, b]$, så gjelder

$$f(x) - p(x) = \frac{1}{(n+1)!} f^{(n+1)}(\xi) \prod_{i=0}^n (x - x_i), \quad \xi \in (a, b).$$

- Newtons dividerte differansers interpolasjonspolynom $p(x)$ av grad $\leq n$:

$$\begin{aligned} p(x) &= f[x_0] + (x - x_0)f[x_0, x_1] + (x - x_0)(x - x_1)f[x_0, x_1, x_2] \\ &\quad + \dots + (x - x_0)(x - x_1)\dots(x - x_{n-1})f[x_0, \dots, x_n] \end{aligned}$$

- Numerisk derivasjon:

$$\begin{aligned} f'(x) &= \frac{1}{h} [f(x+h) - f(x)] + \frac{1}{2} h f''(\xi) \\ f'(x) &= \frac{1}{2h} [f(x+h) - f(x-h)] - \frac{1}{6} h^2 f'''(\xi) \\ f''(x) &= \frac{1}{h^2} [f(x+h) - 2f(x) + f(x-h)] - \frac{1}{12} h^2 f^{(4)}(\xi) \end{aligned}$$

- Simpsons integrasjonsformel:

$$\int_{x_0}^{x_2} f(x) \, dx \approx \frac{h}{3} (f_0 + 4f_1 + f_2)$$

- Newtons metode for likninga $f(x) = 0$ er gitt ved

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

- Iterative teknikkar for løysing av eit lineært likningssystem

$$\sum_{j=1}^n a_{ij}x_j = b_i, \quad i = 1, 2, \dots, n$$

$$\text{Jacobi: } x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij}x_j^{(k)} - \sum_{j=i+1}^n a_{ij}x_j^{(k)} \right)$$

$$\text{Gauss-Seidel: } x_i^{(k+1)} = \frac{1}{a_{ii}} \left(b_i - \sum_{j=1}^{i-1} a_{ij}x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij}x_j^{(k)} \right)$$

- Heuns metode for løysing av $\mathbf{y}' = \mathbf{f}(x, \mathbf{y})$:

$$\begin{aligned} \mathbf{k}_1 &= h\mathbf{f}(x_n, \mathbf{y}_n) \\ \mathbf{k}_2 &= h\mathbf{f}(x_n + h, \mathbf{y}_n + \mathbf{k}_1) \\ \mathbf{y}_{n+1} &= \mathbf{y}_n + \frac{1}{2}(\mathbf{k}_1 + \mathbf{k}_2) \end{aligned}$$

Tabell over nokre Laplace-transformer

| $f(t)$ | $F(s) = \mathcal{L}\{f(t)\} = \int_0^\infty e^{-st} f(t) dt$ |
|--------------------------------|--|
| 1 | $\frac{1}{s}$ |
| t | $\frac{1}{s^2}$ |
| t^n ($n = 0, 1, 2, \dots$) | $\frac{n!}{s^{n+1}}$ |
| e^{at} | $\frac{1}{s - a}$ |
| $\cos \omega t$ | $\frac{s}{s^2 + \omega^2}$ |
| $\sin \omega t$ | $\frac{\omega}{s^2 + \omega^2}$ |
| $\cosh at$ | $\frac{s}{s^2 - a^2}$ |
| $\sinh at$ | $\frac{a}{s^2 - a^2}$ |
| $e^{at} \cos \omega t$ | $\frac{s - a}{(s - a)^2 + \omega^2}$ |
| $e^{at} \sin \omega t$ | $\frac{\omega}{(s - a)^2 + \omega^2}$ |
| $\delta(t - a)$ | e^{-as} |

Tabell over nokre Fourier-transformer

| $f(x)$ | $\widehat{f}(w) = \mathcal{F}\{f(x)\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-iwx} dx$ |
|-------------------|---|
| $g(x) = f(ax)$ | $\widehat{g}(w) = \frac{1}{a} \widehat{f}\left(\frac{w}{a}\right)$ |
| $u(x) - u(x - a)$ | $\frac{1}{\sqrt{2\pi}} \left(\frac{\sin aw}{w} - i \frac{1 - \cos aw}{w} \right)$ |
| e^{-ax^2} | $\frac{1}{\sqrt{2a}} e^{-\frac{w^2}{4a}}$ |