



Norwegian University of Science
and Technology
Department of Mathematical
Sciences

TMA4220 Numerical
Solution of Partial
Differential Equations
Using Element Methods
Fall 2013

Project 2

- 1 If you are not familiar with the the Lebesgue spaces $L^p(\Omega)$ and the Sobolov spaces $H^p(\Omega)$, you should read section 2.3.1 and 2.4.0-2.4.2.

True or false:

- a) The set $S = \{v \in C^0(0,1) : v(\frac{1}{2}) = 1\}$ is a linear (vector) space.
Solution: False. If $v_1, v_2 \in S$, then $v_1 + v_2 \notin S$ since $(v_1 + v_2)(\frac{1}{2}) = 1$.
- b) For $X = H_0^1((0,1))$, $L(v) = \int_0^1 xvdx$ is a linear functional.
Solution: True: $L : X \rightarrow \mathbb{R}$ and $L(\alpha_1v_1 + \alpha_2v_2) = \alpha_1L(v_1) + \alpha_2L(v_2)$, $\forall \alpha_1, \alpha_2 \in \mathbb{R}$ and $v_1, v_2 \in X$.
- c) For $Z = \mathbb{R}$, $(x, y)_Z = |x| |y|$ is a valid inner product.
Solution: False: $(x, y)_Z$ is not bilinear. E.g $(x_1 + x_2, y)_Z = |x_1 + x_2| |y| \neq |x_1| |y| + |x_2| |y|$ if x_1 and x_2 has opposite signs.
- d) The only v in $H^1(\Omega)$ for which $|w|_{H^1(\Omega)}$ (the H^1 semi-norm) is zero is $v = 0$.
Solution: False: Try $w = 1$.
- e) The function $v = x^{3/4}$ is in $L^2((0,1))$; in $H^1((0,1))$; in $H^2((0,1))$.
Solution: We have

$$\begin{aligned}\|v\|_{L^2(\Omega)}^2 &= \int_0^1 (x^{3/2})^2 dx = \frac{2}{3} < \infty \\ \|v_x\|_{L^2(\Omega)}^2 &= \int_0^1 v_x^2 dx = \frac{9}{8} < \infty \\ \|v_{xx}\|_{L^2(\Omega)} &= \int_0^1 v_{xx}^2 dx \not< \infty\end{aligned}$$

So $v \in L_2((0,1))$ and $v \in H^1((0,1))$, but not in $H^2((0,1))$.

- f) For $v = e^{-10x}$, $|v|_{H^2((0,1))} = |v|_{H^1((0,1))}$.
Solution: False.

- 2 Consider the fourth-order problem:

$$\begin{aligned}u_{xxxx} &= f \quad \text{in } \Omega = (0,1), \\ u(0) &= u_x(0) = u(1) = u_x(1) = 0.\end{aligned}$$

This “biharmonic” equation is relevant to, amongst other applications, the bending of beams.

- a) Find a symmetric, positive form a over V and a linear form F such that the solution u of the equation satisfies

$$a(u, v) = F(v), \quad \forall v \in V.$$

Solution: As always, choose a test function $v \in V$, integrate over the domain, and apply partial integration:

$$\int_0^1 u_{xxxx} v dx = u_{xxx} v \Big|_0^1 - \int_0^1 u_{xxx} v_x dx = u_{xxx} v \Big|_0^1 - u_{xx} v_x \Big|_0^1 + \int_0^1 u_{xx} v_{xx} dx = \int_0^1 f v dx.$$

The two boundary terms disappear if $v(0) = v(1) = v_x(0) = v_x(1) = 0$, so we can use $a(u, v) = \int_0^1 u_{xx} v_{xx} dx$ and $F(v) = \int_0^1 f v dx$. Further, a is clearly symmetric, and it is positive definite since $a(v, v) = \int_0^1 v_{xx}^2 dx \geq 0$, and $a(v, v) = 0$ only if $v_{xx} \equiv 0$ which implies $v \equiv 0$ because of the boundary conditions.

- b) How should V be defined?

Solution: $V = H_0^2((0, 1)) = \{v \in H^2((0, 1)) : v(0) = v(1) = v_x(0) = v_x(1) = 0\}$.

- c) Do you think that $F(v) = v_x(\frac{1}{2})$ is a linear, bounded functional on V ?

Solution: Yes, since

$$\begin{aligned} F(v) &= v_x\left(\frac{1}{2}\right) = \int_0^{1/2} v_{xx}(x) dx && \text{since } v_x(0) = 0 \\ |F(v)| &\leq \int_0^{1/2} |v_{xx}(x)| \cdot 1 dx \leq \left(\int_0^{1/2} v_{xx}^2 dx\right)^{1/2} \left(\int_0^{1/2} 1^2 dx\right)^{1/2} && \text{Cauchy-Schwartz} \\ &= \frac{\sqrt{2}}{2} \left(\int_0^{1/2} v_{xx}^2 dx\right)^{1/2} \leq \frac{\sqrt{2}}{2} \left(\int_0^1 v_{xx}^2 dx\right)^{1/2} \\ &= \frac{\sqrt{2}}{2} \|v\|_{H^2((0,1))} \leq \frac{\sqrt{2}}{2} \|v\|_{H^2((0,1))}. \end{aligned}$$

3 Consider the problem with a discontinuous jump in conductivities:

$$\begin{aligned} -\kappa^L u_{xx}^L &= f^L, & 0 < x < \frac{1}{2}, \\ -\kappa^R u_{xx}^R &= f^R, & \frac{1}{2} < x < 1, \end{aligned}$$

with boundary conditions

$$\begin{aligned} u^L(0) &= 0, & u^R(1) &= 0, \\ u^L\left(\frac{1}{2}\right) &= u^R\left(\frac{1}{2}\right), \\ \kappa^L u_x^L\left(\frac{1}{2}\right) &= \kappa^R u_x^R\left(\frac{1}{2}\right), & (\text{continuity of flux}). \end{aligned}$$

Here, κ^L and κ^R are strictly positive. Let $V = \{v \in H^1((0, 1)) : v(0) = v(1) = 0\}$. Find a and F such that the solution u satisfies

$$a(u, v) = F(v), \quad \forall v \in V.$$

Solution: Choose some $v \in V$. Then

$$\begin{aligned} -\int_0^{1/2} \kappa^L u_{xx}^L v dx - \int_{1/2}^1 \kappa^R u_{xx}^R v dx &= -\kappa^L u_x v \Big|_0^{1/2} + \int_0^{1/2} \kappa^L u_x^L v_x dx - \kappa^R u_x^R v \Big|_{1/2}^1 + \int_{1/2}^1 \kappa^R u_x^R v dx \\ &= \left(-\kappa^L u_x^L \left(\frac{1}{2} \right) + \kappa^R u_x^R \left(\frac{1}{2} \right) \right) v \left(\frac{1}{2} \right) + \int_0^{1/2} \kappa^L u_x^L v_x dx + \int_{1/2}^1 \kappa^R u_x^R v_x dx \\ &= \int_0^{1/2} f^L v dx + \int_{1/2}^1 f^R v dx \end{aligned}$$

The first term of the second line is zero because of the continuity of flux condition. So we can choose

$$a(u, v) = \int_0^{1/2} \kappa^L u_x^L v_x dx + \int_{1/2}^1 \kappa^R u_x^R v_x dx, \quad F(v) = \int_0^{1/2} f^L v dx + \int_{1/2}^1 f^R v dx.$$

4 Given the Helmholtz problem

$$\begin{aligned} -u_{xx} + \sigma u &= f \text{ on } (0, 1), \\ u(0) &= u(1) = 0. \end{aligned}$$

where $\sigma > 0$ is a constant. Set up the weak form for this problem. Show that, when this problem is solved by a Galerkin method, using $V_h = \text{span} \{ \phi_i \}_{i=1}^N$, the discrete problem can be written as

$$(A + \sigma M)\mathbf{u} = \mathbf{f}.$$

Set up the matrix M for $V_h = X_h^1$ on a uniform grid.

Solution: Multiply by the equation by a test function v , integrate over the domain $(0, 1)$, use partial integration and get rid of the boundary terms by require $v(0) = v(1) = 0$. The the weak formulation becomes:

$$\text{Find } u \in H_0^1(\Omega) \text{ such that } \int_0^1 v_x u_x dx + \sigma \int_0^1 u v dx = \int_0^1 f v dx, \quad \forall v \in H_0^1(0, 1).$$

Choose $V_h = \text{span}\{\varphi_1, \varphi_2, \dots, \varphi_N\}$, let our unknown approximation be written as

$$u_h(x) = \sum_{j=1}^N u_j \varphi_j(x),$$

where the coefficients u_j is found from

$$\sum_{j=1}^N \int_0^1 \left(\frac{d\varphi_j}{dx} \frac{d\varphi_i}{dx} \right) dx + \sigma \sum_{j=1}^N \int_0^1 (\varphi_j \varphi_i) dx = \int_0^1 (f \varphi_i) dx, \quad i = 1, 2, \dots, N.$$


This is a linear system of equations

$$A\mathbf{u} + \sigma M\mathbf{u} = \mathbf{f}$$

where

$$(A_{i,j} = \int_0^1 \left(\frac{d\varphi_j}{dx} \frac{d\varphi_i}{dx} \right) dx, \quad M_{i,j} = \int_0^1 (\varphi_j \varphi_i) dx \quad \text{and} \quad f_i = \int_0^1 (f \varphi_i) dx.$$

Let $V_h = X_h^1$ on a uniform grid, so that $h = 1/N$ and $x_i = ih$, $i = 0, 1, \dots, N$.



The non-zero elements of the stiffness matrix A and the mass matrix M is

$$A_{i,i} = \int_{x_{i-1}}^{x_{i+1}} \left(\frac{d\varphi_i}{dx} \right) dx = \frac{2}{h}, \quad A_{i,i+1} = \int_{x_i}^{x_{i+1}} \frac{d\varphi_{i+1}}{dx} \frac{d\varphi_i}{dx} dx = -\frac{1}{h} = A_{i+1,i}$$

$$M_{i,i} = \int_{x_{i-1}}^{x_{i+1}} \varphi_i^2 dx = \frac{2}{3}h, \quad M_{i,i+1} = \int_{x_i}^{x_{i+1}} \varphi_{i+1} \varphi_i dx = \frac{1}{6}h = M_{i+1,i}.$$

All the other elements are 0. So

$$M = \frac{h}{6} \begin{pmatrix} 4 & 1 & & & 0 \\ 1 & 4 & 1 & & \\ & 1 & 4 & \ddots & \\ & & \ddots & \ddots & 1 \\ 0 & & & 1 & 4 \end{pmatrix} \in \mathbb{R}^{(N-1) \times (N-1)}.$$