

THE AXIOMS FOR n -ANGULATED CATEGORIES

PETTER ANDREAS BERGH AND MARIUS THAULE

ABSTRACT. We discuss the axioms for an n -angulated category, recently introduced by Geiss, Keller and Oppermann in [2]. In particular, we introduce a higher “octahedral axiom”, and show that it is equivalent to the mapping cone axiom for an n -angulated category. For a triangulated category, the mapping cone axiom, our octahedral axiom and the classical octahedral axiom are all equivalent.

1. INTRODUCTION

Triangulated categories were introduced independently in algebraic geometry by Verdier [9, 10], based on ideas of Grothendieck, and in algebraic topology by Puppe [8]. These constructions have since played a crucial role in representation theory, algebraic geometry, commutative algebra, algebraic topology and other areas of mathematics (and even theoretical physics). Recently, Geiss, Keller and Oppermann introduced in [2] a new type of categories, called n -angulated categories, which generalize triangulated categories: the classical triangulated categories are the special case $n = 3$. These categories appear for instance when considering certain $(n - 2)$ -cluster tilting subcategories of triangulated categories. Conversely, certain n -angulated Calabi–Yau categories yield triangulated Calabi–Yau categories of higher Calabi–Yau dimension.

The four axioms for n -angulated categories are generalizations of the axioms for triangulated categories. In this paper, we discuss these axioms, inspired by works of Neeman [6, 7]. First, we show that the first two of the original axioms can be replaced by two alternative axioms. One of these alternative axioms requires that the collection of n -angles be closed under so-called weak isomorphisms, but not under direct sums and summands. The other axiom requires that the collection of n -angles be closed only under left rotations, but not right rotations. Second, we discuss the axioms that enable us to complete certain diagrams to morphisms of n -angles. The last of these axioms says that we can complete diagrams to morphisms of n -angles in such a way that the mapping cone is itself an n -angle. For triangulated categories (that is, when $n = 3$), this axiom is equivalent to the octahedral axiom, which was one of Verdier’s original axioms. We show that this generalizes to n -angulated categories. Namely, we introduce a higher “octahedral axiom” for n -angulated categories, and show that this is equivalent to the mapping cone axiom. For $n = 3$, that is, for triangulated categories, our new axiom is almost the same as the classical octahedral axiom. In fact, it is apparently a bit weaker, but we show that they are equivalent. Therefore, for a triangulated category, the mapping cone axiom, our octahedral axiom and the classical octahedral axiom are all equivalent.

This paper is organized as follows. In Section 2, we recall the definition of n -angulated categories from [2]. Then, in Section 3, we discuss the first two axioms, and in Section 4 we introduce the higher octahedral axiom. Finally, in Section 5, we look at an example, namely the n -angulated categories originating from $(n - 2)$ -cluster tilting subcategories of triangulated categories. We verify the higher octahedral axiom for these categories, in the case when $n = 4$.

2. THE AXIOMS FOR n -ANGULATED CATEGORIES

Throughout this paper, we fix an additive category \mathcal{C} with an automorphism $\Sigma: \mathcal{C} \rightarrow \mathcal{C}$, and an integer n greater than or equal to three. In this section, we recall the set of axioms for n -angulated categories as described in [2].

A sequence of objects and morphisms in \mathcal{C} of the form

$$A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} \dots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma A_1$$

2010 *Mathematics Subject Classification.* 18E30.

Key words and phrases. Triangulated categories, n -angulated categories, octahedral axiom.

The second author was supported by the Research Council of Norway grant Topology, grant number 185335/V30.

is called an n - Σ -sequence; we shall frequently denote such sequences by A_\bullet, B_\bullet etc. The n - Σ -sequence A_\bullet is *exact* if the induced sequence

$$\cdots \rightarrow \mathrm{Hom}_{\mathcal{C}}(B, A_1) \xrightarrow{(\alpha_1)_*} \mathrm{Hom}_{\mathcal{C}}(B, A_2) \xrightarrow{(\alpha_2)_*} \cdots \xrightarrow{(\alpha_{n-1})_*} \mathrm{Hom}_{\mathcal{C}}(B, A_n) \xrightarrow{(\alpha_n)_*} \mathrm{Hom}_{\mathcal{C}}(B, \Sigma A_1) \rightarrow \cdots$$

of abelian groups is exact for every object $B \in \mathcal{C}$. The left and right *rotations* of A_\bullet are the two n - Σ -sequences

$$A_2 \xrightarrow{\alpha_2} A_3 \xrightarrow{\alpha_3} \cdots \xrightarrow{\alpha_n} \Sigma A_1 \xrightarrow{(-1)^n \Sigma \alpha_1} \Sigma A_2$$

and

$$\Sigma^{-1} A_n \xrightarrow{(-1)^n \Sigma^{-1} \alpha_n} A_1 \xrightarrow{\alpha_1} \cdots \xrightarrow{\alpha_{n-2}} A_{n-1} \xrightarrow{\alpha_{n-1}} A_n$$

respectively, and a *trivial* n - Σ -sequence is a sequence of the form

$$A \xrightarrow{1} A \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \Sigma A$$

or any of its rotations.

A *morphism* $A_\bullet \xrightarrow{\varphi} B_\bullet$ of n - Σ -sequences is a sequence $\varphi = (\varphi_1, \varphi_2, \dots, \varphi_n)$ of morphisms in \mathcal{C} such that the diagram

$$\begin{array}{ccccccccccc} A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\ \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma \varphi_1 \\ B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1 \end{array}$$

commutes. It is an *isomorphism* if $\varphi_1, \varphi_2, \dots, \varphi_n$ are all isomorphisms in \mathcal{C} , and a *weak isomorphism* if φ_i and φ_{i+1} are isomorphisms for some $1 \leq i \leq n$ (with $\varphi_{n+1} := \Sigma \varphi_1$). Note that the composition of two weak isomorphisms need not be a weak isomorphism. Also, note that if two n - Σ -sequences A_\bullet and B_\bullet are weakly isomorphic through a weak isomorphism $A_\bullet \xrightarrow{\varphi} B_\bullet$, then there does not necessarily exist a weak isomorphism $B_\bullet \rightarrow A_\bullet$ in the opposite direction.

The category \mathcal{C} is *pre- n -angulated* if there exists a collection \mathcal{N} of n - Σ -sequences satisfying the following three axioms:

- (N1) (a) \mathcal{N} is closed under direct sums, direct summands and isomorphisms of n - Σ -sequences.
(b) For all $A \in \mathcal{C}$, the trivial n - Σ -sequence

$$A \xrightarrow{1} A \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \Sigma A$$

belongs to \mathcal{N} .

- (c) For each morphism $\alpha: A_1 \rightarrow A_2$ in \mathcal{C} , there exists an n - Σ -sequence in \mathcal{N} whose first morphism is α .
(N2) An n - Σ -sequence belongs to \mathcal{N} if and only if its left rotation belongs to \mathcal{N} .
(N3) Each commutative diagram

$$\begin{array}{ccccccccccc} A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\ \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma \varphi_1 \\ B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1 \end{array}$$

with rows in \mathcal{N} can be completed to a morphism of n - Σ -sequences.

In this case, the collection \mathcal{N} is a *pre- n -angulation* of the category \mathcal{C} (relative to the automorphism Σ), and the n - Σ -sequences in \mathcal{N} are *n -angles*. If, in addition, the collection \mathcal{N} satisfies the following axiom, then it is an *n -angulation* of \mathcal{C} , and the category is *n -angulated*:

- (N4) In the situation of (N3), the morphisms $\varphi_3, \varphi_4, \dots, \varphi_n$ can be chosen such that the mapping cone

$$A_2 \oplus B_1 \xrightarrow{\begin{bmatrix} -\alpha_2 & 0 \\ \varphi_2 & \beta_1 \end{bmatrix}} A_3 \oplus B_2 \xrightarrow{\begin{bmatrix} -\alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} \cdots \xrightarrow{\begin{bmatrix} -\alpha_n & 0 \\ \varphi_n & \beta_{n-1} \end{bmatrix}} \Sigma A_1 \oplus B_n \xrightarrow{\begin{bmatrix} -\Sigma \alpha_1 & 0 \\ \Sigma \varphi_1 & \beta_n \end{bmatrix}} \Sigma A_2 \oplus \Sigma B_1$$

belongs to \mathcal{N} .

Note that in [2], it was not explicitly assumed that \mathcal{N} be closed under isomorphisms, but it follows implicitly from closure under direct sums. Since closure under isomorphisms is a crucial part of many of our proofs, we have included it as a part of axiom (N1)(a). Note also that by [2, Proposition 1.5], every n -angle in a pre- n -angulated category is exact. Consequently, the composition of two consecutive morphisms in an

n -angle is zero. Finally, note the similarity with Balmer's recent definition (cf. [1]) of triangulated categories of order n .

3. AXIOMS (N1) AND (N2)

In this section, we discuss the first two defining axioms (N1) and (N2) for pre- n -angulated categories. It turns out that we may replace these axioms by the following ones:

- (N1*) (a) If $A_\bullet \xrightarrow{\varphi} B_\bullet$ is a weak isomorphism of exact n - Σ -sequences with $A_\bullet \in \mathcal{N}$, then B_\bullet belongs to \mathcal{N} .
 (b) For all $A \in \mathcal{C}$, the trivial n - Σ -sequence

$$A \xrightarrow{1} A \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \Sigma A$$

belongs to \mathcal{N} .

- (c) For each morphism $\alpha: A_1 \rightarrow A_2$ in \mathcal{C} , there exists an n - Σ -sequence in \mathcal{N} whose first morphism is α .

- (N2*) The left rotation of every n - Σ -sequence in \mathcal{N} also belongs to \mathcal{N} .

In axiom (N1*), we do not require that \mathcal{N} be closed under direct sums and summands. However, we do require that \mathcal{N} be closed under weak isomorphisms (in one direction), and this is stronger than requiring that \mathcal{N} be closed under isomorphisms. In axiom (N2*), we only require that \mathcal{N} be closed under left rotations. This is sometimes done when considering triangulated categories, cf. [4].

Because of the new axiom (N1*)(a), the exact n - Σ -sequences play an important role in the proofs to come. We therefore need to determine which properties a collection \mathcal{N} of n - Σ -sequences must satisfy in order for all its elements to be exact. We do this in the following result.

Lemma 3.1. *If \mathcal{N} is a collection of n - Σ -sequences satisfying the axioms (N1)(b), (N2*) and (N3), then all the elements in \mathcal{N} are exact.*

Proof. Let

$$A_\bullet: \quad A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma A_1$$

be an n - Σ -sequence in \mathcal{N} , and pick an integer $1 \leq j \leq n$. In the diagram

$$\begin{array}{ccccccccccc} A_j & \xrightarrow{1} & A_j & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & 0 & \longrightarrow & \Sigma A_j \\ \parallel & & \downarrow \alpha_j & & \vdots & & & & \vdots & & \parallel \\ A_j & \xrightarrow{\alpha_j} & A_{j+1} & \xrightarrow{\alpha_{j+1}} & A_{j+2} & \xrightarrow{\alpha_{j+2}} & \cdots & \xrightarrow{(-1)^n \Sigma \alpha_{j-2}} & \Sigma A_{j-1} & \xrightarrow{(-1)^n \Sigma \alpha_{j-1}} & \Sigma A_j \end{array}$$

the two rows both belong to \mathcal{N} : the top row by (N1)(b), and the bottom row by (repeated use of) (N2*). If $j = 1$ or $j = 2$, the two rightmost morphisms in the bottom row have different labels, and if $j = n$ then $\alpha_{n+1} = (-1)^n \Sigma \alpha_1$. By (N3), we can complete the diagram to a morphism of n - Σ -sequences, hence

$$\alpha_2 \circ \alpha_1 = \alpha_3 \circ \alpha_2 = \cdots = \alpha_n \circ \alpha_{n-1} = (\Sigma \alpha_1) \circ \alpha_n = 0.$$

For objects $X, Y \in \mathcal{C}$, denote the abelian group $\text{Hom}_{\mathcal{C}}(X, Y)$ by (X, Y) . Since all the possible compositions of morphisms from A_\bullet are zero, the doubly infinite sequence

$$\cdots \rightarrow (B, \Sigma^{i-1} A_n) \xrightarrow{(\Sigma^{i-1} \alpha_n)_*} (B, \Sigma^i A_1) \xrightarrow{(\Sigma^i \alpha_1)_*} \cdots \xrightarrow{(\Sigma^i \alpha_{n-1})_*} (B, \Sigma^i A_n) \xrightarrow{(\Sigma^i \alpha_n)_*} (B, \Sigma^{i+1} A_1) \rightarrow \cdots$$

of abelian groups and maps is a complex for every object $B \in \mathcal{C}$. Now pick an integer $1 \leq i \leq n$, and let f be an element in $\text{Ker}(\Sigma^i \alpha_j)_*$. Then f is a morphism in $\text{Hom}_{\mathcal{C}}(B, \Sigma^i A_j)$ with $(\Sigma^i \alpha_j) \circ f = 0$. Applying the automorphism Σ^{-i} , we obtain $\alpha_j \circ (\Sigma^{-i} f) = 0$, where $\Sigma^{-i} f$ is a morphism in $\text{Hom}_{\mathcal{C}}(\Sigma^{-i} B, A_j)$. Now consider the diagram

$$\begin{array}{ccccccccccc} \Sigma^{-i} B & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & \Sigma^{1-i} B & \xrightarrow{(-1)^n} & \Sigma^{1-i} B \\ \downarrow \Sigma^{-i} f & & \downarrow & & & & \downarrow g & & \downarrow \Sigma^{1-i} f \\ A_j & \xrightarrow{\alpha_j} & A_{j+1} & \xrightarrow{\alpha_{j+1}} & \cdots & \xrightarrow{(-1)^n \Sigma \alpha_{j-2}} & \Sigma A_{j-1} & \xrightarrow{(-1)^n \Sigma \alpha_{j-1}} & \Sigma A_j \end{array}$$

in which the two rows belong to \mathcal{N} by (N1)(b) and (repeated use of) (N2*). By (N3), we can complete this diagram to a morphism of n - Σ -sequences, and in particular we obtain a morphism $g \in \text{Hom}_{\mathcal{C}}(\Sigma^{1-i}B, \Sigma A_{j-1})$ with

$$(\Sigma \alpha_{j-1}) \circ g = \Sigma^{1-i} f.$$

Applying the automorphism Σ^{i-1} gives

$$f = (\Sigma^i \alpha_{j-1}) \circ (\Sigma^{i-1} g),$$

hence $f \in \text{Im}(\Sigma^i \alpha_{j-1})_*$. This shows that the complex is exact, and so A_\bullet is an exact n - Σ -sequence. \square

We may now prove that axiom (N1) can be replaced with axiom (N1*).

Theorem 3.2. *If \mathcal{N} is a collection of n - Σ -sequences satisfying the axioms (N2) and (N3), then the following are equivalent:*

- (1) \mathcal{N} satisfies (N1),
- (2) \mathcal{N} satisfies (N1*).

Proof. The implication (1) \Rightarrow (2) is part of [2, Lemma 1.4], hence we must prove that (1) follows from (2), i.e. that \mathcal{N} satisfies (N1)(a) whenever it satisfies (N1*). Suppose therefore that \mathcal{N} satisfies (N1*).

Since the collection \mathcal{N} satisfies the axioms (N1*)(b), (N2) and (N3), the n - Σ -sequences in \mathcal{N} are exact by Lemma 3.1. Now let A_\bullet and B_\bullet be isomorphic n - Σ -sequences, with A_\bullet in \mathcal{N} . Then A_\bullet is exact, and so B_\bullet must also be exact since it is isomorphic to A_\bullet . Since A_\bullet and B_\bullet are trivially weakly isomorphic through an isomorphism $A_\bullet \rightarrow B_\bullet$, the n - Σ -sequence B_\bullet also belongs to \mathcal{N} . This shows that \mathcal{N} is closed under isomorphisms.

Next, we show that \mathcal{N} is closed under direct sums. Given two n - Σ -sequences

$$\begin{array}{l} A_\bullet : \quad A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma A_1 \\ B_\bullet : \quad B_1 \xrightarrow{\beta_1} B_2 \xrightarrow{\beta_2} \cdots \xrightarrow{\beta_{n-1}} B_n \xrightarrow{\beta_n} \Sigma B_1 \end{array}$$

in \mathcal{N} , the direct sum $A_\bullet \oplus B_\bullet$ is exact, since each of the sequences is exact by the above. Now use (N1*)(c) to complete the first morphism in $A_\bullet \oplus B_\bullet$ to an n - Σ -sequence

$$A_1 \oplus B_1 \xrightarrow{\begin{bmatrix} \alpha_1 & 0 \\ 0 & \beta_1 \end{bmatrix}} A_2 \oplus B_2 \xrightarrow{\gamma_2} C_3 \xrightarrow{\gamma_3} \cdots \xrightarrow{\gamma_{n-1}} C_n \xrightarrow{\gamma_n} \Sigma A_1 \oplus \Sigma B_1$$

in \mathcal{N} . By (N3), the two commutative diagrams

$$\begin{array}{ccccccccccc} A_1 \oplus B_1 & \xrightarrow{\begin{bmatrix} \alpha_1 & 0 \\ 0 & \beta_1 \end{bmatrix}} & A_2 \oplus B_2 & \xrightarrow{\gamma_2} & C_3 & \xrightarrow{\gamma_3} & \cdots & \xrightarrow{\gamma_{n-1}} & C_n & \xrightarrow{\gamma_n} & \Sigma A_1 \oplus \Sigma B_1 \\ \downarrow [1 \ 0] & & \downarrow [1 \ 0] & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow [1 \ 0] \\ A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\ \\ A_1 \oplus B_1 & \xrightarrow{\begin{bmatrix} \alpha_1 & 0 \\ 0 & \beta_1 \end{bmatrix}} & A_2 \oplus B_2 & \xrightarrow{\gamma_2} & C_3 & \xrightarrow{\gamma_3} & \cdots & \xrightarrow{\gamma_{n-1}} & C_n & \xrightarrow{\gamma_n} & \Sigma A_1 \oplus \Sigma B_1 \\ \downarrow [0 \ 1] & & \downarrow [0 \ 1] & & \downarrow \psi_3 & & & & \downarrow \psi_n & & \downarrow [0 \ 1] \\ B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1 \end{array}$$

can be completed to morphisms of n - Σ -sequences, since the sequences involved are all in \mathcal{N} . This gives a weak isomorphism

$$\begin{array}{ccccccccccc} A_1 \oplus B_1 & \xrightarrow{\begin{bmatrix} \alpha_1 & 0 \\ 0 & \beta_1 \end{bmatrix}} & A_2 \oplus B_2 & \xrightarrow{\gamma_2} & C_3 & \xrightarrow{\gamma_3} & \cdots & \xrightarrow{\gamma_{n-1}} & C_n & \xrightarrow{\gamma_n} & \Sigma A_1 \oplus \Sigma B_1 \\ \parallel & & \parallel & & \downarrow \begin{bmatrix} \varphi_3 \\ \psi_3 \end{bmatrix} & & & & \downarrow \begin{bmatrix} \varphi_n \\ \psi_n \end{bmatrix} & & \parallel \\ A_1 \oplus B_1 & \xrightarrow{\begin{bmatrix} \alpha_1 & 0 \\ 0 & \beta_1 \end{bmatrix}} & A_2 \oplus B_2 & \xrightarrow{\begin{bmatrix} \alpha_2 & 0 \\ 0 & \beta_2 \end{bmatrix}} & A_3 \oplus B_3 & \xrightarrow{\begin{bmatrix} \alpha_3 & 0 \\ 0 & \beta_3 \end{bmatrix}} & \cdots & \xrightarrow{\begin{bmatrix} \alpha_{n-1} & 0 \\ 0 & \beta_{n-1} \end{bmatrix}} & A_n \oplus B_n & \xrightarrow{\begin{bmatrix} \alpha_n & 0 \\ 0 & \beta_n \end{bmatrix}} & \Sigma A_1 \oplus \Sigma B_1 \end{array}$$

of n - Σ -sequences. The top sequence belongs to \mathcal{N} and is therefore exact, whereas the bottom sequence $A_\bullet \oplus B_\bullet$ is also exact. From (N1*)(a) we conclude that $A_\bullet \oplus B_\bullet$ belongs to \mathcal{N} .

Finally, we show that \mathcal{N} is closed under direct summands. Suppose therefore that A_\bullet and B_\bullet are n - Σ -sequences as above, that B_\bullet belongs to \mathcal{N} (hence B_\bullet is exact), and that A_\bullet is a direct summand of B_\bullet . Then there exists a diagram

$$\begin{array}{ccccccccccc}
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
 \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma \varphi_1 \\
 B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1 \\
 \downarrow \psi_1 & & \downarrow \psi_2 & & \downarrow \psi_3 & & & & \downarrow \psi_n & & \downarrow \Sigma \psi_1 \\
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1
 \end{array}$$

of morphisms $A_\bullet \xrightarrow{\varphi} B_\bullet$ and $B_\bullet \xrightarrow{\psi} A_\bullet$ of n - Σ -sequences, with $\psi_i \circ \varphi_i = 1_{A_i}$ for all i . For every object Z in \mathcal{C} , the sequence $\text{Hom}_{\mathcal{C}}(Z, A_\bullet)$ of abelian groups and maps is a direct summand of the exact sequence $\text{Hom}_{\mathcal{C}}(Z, B_\bullet)$, and is therefore itself exact. Consequently, the n - Σ -sequence A_\bullet is exact. Now use (N1*)(c) to complete the first morphism in A_\bullet to an n - Σ -sequence

$$D_\bullet: \quad A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\delta_2} D_3 \xrightarrow{\delta_3} \cdots \xrightarrow{\delta_{n-1}} D_n \xrightarrow{\delta_n} \Sigma A_1$$

in \mathcal{N} (in particular, D_\bullet is exact). Using this sequence, we can obtain a diagram

$$\begin{array}{ccccccccccc}
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\delta_2} & D_3 & \xrightarrow{\delta_3} & \cdots & \xrightarrow{\delta_{n-1}} & D_n & \xrightarrow{\delta_n} & \Sigma A_1 \\
 \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \theta_3 & & & & \downarrow \theta_n & & \downarrow \Sigma \varphi_1 \\
 B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1 \\
 \downarrow \psi_1 & & \downarrow \psi_2 & & \downarrow \psi_3 & & & & \downarrow \psi_n & & \downarrow \Sigma \psi_1 \\
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1
 \end{array}$$

whose rows are D_\bullet , B_\bullet and A_\bullet . The top half of this diagram is a morphism $D_\bullet \xrightarrow{\theta} B_\bullet$, which we obtain from (N3), whereas the lower half is the morphism $B_\bullet \xrightarrow{\psi} A_\bullet$. Moreover, the composition $D_\bullet \xrightarrow{\psi \circ \theta} A_\bullet$ is a weak isomorphism, since $\psi_1 \circ \varphi_1 = 1_{A_1}$ and $\psi_2 \circ \varphi_2 = 1_{A_2}$. Since both D_\bullet and A_\bullet are exact, and $D_\bullet \in \mathcal{N}$, the sequence A_\bullet belongs to \mathcal{N} by (N1*)(a). This shows that the collection \mathcal{N} is closed under direct summands. We have now proved that \mathcal{N} is closed under isomorphisms, direct sums and direct summands, which is axiom (N1)(a). \square

Next, we study the rotation axiom (N2). The following result shows that when we replace (N1) with (N1*), then we can also replace (N2) with the weaker version with (N2*). In other words, in the rotation axiom we only need to require that the left rotation of an n - Σ sequence in \mathcal{N} also belongs to \mathcal{N} .

Theorem 3.3. *If \mathcal{N} is a collection of n - Σ -sequences satisfying the axioms (N1*) and (N3), then the following are equivalent:*

- (1) \mathcal{N} satisfies (N2),
- (2) \mathcal{N} satisfies (N2*).

Proof. The implication (1) \Rightarrow (2) is trivial. Assume therefore that \mathcal{N} satisfies (N2*), and let

$$A_\bullet: \quad A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma A_1$$

be an n - Σ -sequence in \mathcal{N} . By repeatedly applying (N2*), we obtain the n - Σ -sequence

$$A_n \xrightarrow{\alpha_n} \Sigma A_1 \xrightarrow{(-1)^n \Sigma \alpha_1} \cdots \xrightarrow{(-1)^n \Sigma \alpha_{n-2}} \Sigma A_{n-1} \xrightarrow{(-1)^n \Sigma \alpha_{n-1}} \Sigma A_n$$

in \mathcal{N} . Now use (N1*)(c) to complete the morphism $\Sigma^{-1} A_n \xrightarrow{(-1)^n \Sigma^{-1} \alpha_n} A_1$ to an n - Σ -sequence

$$\Sigma^{-1}A_n \xrightarrow{(-1)^n \Sigma^{-1} \alpha_n} A_1 \xrightarrow{\beta_2} B_3 \xrightarrow{\beta_3} \cdots \xrightarrow{\beta_{n-1}} B_n \xrightarrow{\beta_n} A_n$$

in \mathcal{N} . By repeated use of (N2*), we obtain the n - Σ -sequence

$$A_n \xrightarrow{\alpha_n} \Sigma A_1 \xrightarrow{(-1)^n \Sigma \beta_2} \Sigma B_3 \xrightarrow{(-1)^n \Sigma \beta_3} \cdots \xrightarrow{(-1)^n \Sigma \beta_{n-1}} \Sigma B_n \xrightarrow{(-1)^n \Sigma \beta_n} \Sigma A_n$$

in \mathcal{N} . By (N3), we may complete the diagram

$$\begin{array}{ccccccccccc} A_n & \xrightarrow{\alpha_n} & \Sigma A_1 & \xrightarrow{(-1)^n \Sigma \beta_2} & \Sigma B_3 & \xrightarrow{(-1)^n \Sigma \beta_3} & \cdots & \xrightarrow{(-1)^n \Sigma \beta_{n-1}} & \Sigma B_n & \xrightarrow{(-1)^n \Sigma \beta_n} & \Sigma A_n \\ \parallel & & \parallel & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \parallel \\ A_n & \xrightarrow{\alpha_n} & \Sigma A_1 & \xrightarrow{(-1)^n \Sigma \alpha_1} & \Sigma A_2 & \xrightarrow{(-1)^n \Sigma \alpha_2} & \cdots & \xrightarrow{(-1)^n \Sigma \alpha_{n-2}} & \Sigma A_{n-1} & \xrightarrow{(-1)^n \Sigma \alpha_{n-1}} & \Sigma A_n \end{array}$$

and obtain a morphism of n - Σ -sequences. By applying the automorphism Σ^{-1} to the whole diagram, and multiplying all maps with $(-1)^n$, we obtain a weak isomorphism

$$\begin{array}{ccccccccccc} \Sigma^{-1}A_n & \xrightarrow{(-1)^n \Sigma^{-1} \alpha_n} & A_1 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & A_n \\ \downarrow (-1)^n & & \downarrow (-1)^n & & \downarrow (-1)^n \Sigma^{-1} \varphi_3 & & & & \downarrow (-1)^n \Sigma^{-1} \varphi_n & & \downarrow (-1)^n \\ \Sigma^{-1}A_n & \xrightarrow{(-1)^n \Sigma^{-1} \alpha_n} & A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & \cdots & \xrightarrow{\alpha_{n-2}} & A_{n-1} & \xrightarrow{\alpha_{n-1}} & A_n \end{array}$$

of n - Σ -sequences. The top row belongs to \mathcal{N} and is therefore exact by Lemma 3.1, whereas the bottom row is the right rotation of A_\bullet . Since A_\bullet is exact, so is its right rotation, and from (N1*)(a) we conclude that this right rotation also belongs to \mathcal{N} . \square

Collecting the results in this section gives the following.

Theorem 3.4. *For a collection \mathcal{N} of n - Σ -sequences, the following are equivalent:*

- (1) \mathcal{N} satisfies (N1), (N2) and (N3),
- (2) \mathcal{N} satisfies (N1*), (N2) and (N3),
- (3) \mathcal{N} satisfies (N1*), (N2*) and (N3).

4. AXIOM (N4)

For triangulated categories, it is a well-known fact that Verdier's original octahedral axiom has several equivalent representations, see e.g. [3] for a discussion. It is natural to ask whether this also holds true for general n -angulated categories. We prove in this section that it does: we introduce a higher ‘‘octahedral axiom’’ (N4*) for n -angulated categories, and show that it is equivalent to axiom (N4).

What is the essence of the classical octahedral axiom for triangulated categories? It starts with three given triangles

$$\begin{array}{l} A_1 \rightarrow A_2 \rightarrow A_3 \rightarrow \Sigma A_1 \\ A_1 \rightarrow B_2 \rightarrow B_3 \rightarrow \Sigma A_1 \\ A_2 \rightarrow B_2 \rightarrow C_3 \rightarrow \Sigma A_2 \end{array}$$

that are connected, in that each pair of triangles share a common object. The axiom then guarantees the existence of two new morphisms, and from these new morphisms we obtain three things:

- (1) A morphism of triangles.
- (2) A new triangle, whose objects are objects in the three original triangles.
- (3) Commutativity relations between morphisms.

The reason why the axiom is called the ‘‘octahedral axiom’’ is that everything fits into an octahedron whose vertices are the objects, and where the edges are the morphisms.

The essence of the higher octahedral axiom for n -angulated categories that we now introduce is exactly the same. It starts with three given n -angles, and guarantees the existence of $3n - 7$ new morphisms. From these new morphisms we obtain a morphism of n -angles, a new n -angle and a certain commutativity relation between morphisms.

(N4*) Given a commutative diagram

$$\begin{array}{ccccccccccccccc}
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-2}} & A_{n-1} & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
 \parallel & & \downarrow \varphi_2 & & & & & & & & & & \parallel \\
 A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-2}} & B_{n-1} & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma A_1 \\
 & & \downarrow \gamma_2 & & & & & & & & & & \\
 & & C_3 & & & & & & & & & & \\
 & & \downarrow \gamma_3 & & & & & & & & & & \\
 & & \vdots & & & & & & & & & & \\
 & & \downarrow \gamma_{n-2} & & & & & & & & & & \\
 & & C_{n-1} & & & & & & & & & & \\
 & & \downarrow \gamma_{n-1} & & & & & & & & & & \\
 & & C_n & & & & & & & & & & \\
 & & \downarrow \gamma_n & & & & & & & & & & \\
 & & \Sigma A_2 & & & & & & & & & &
 \end{array}$$

whose top rows and second column are n -angles. Then there exist morphisms $A_i \xrightarrow{\varphi_i} B_i$ ($3 \leq i \leq n$) and ψ_j ($1 \leq j \leq 2n-5$) with the following two properties:

- (1) The sequence $(1, \varphi_2, \varphi_3, \dots, \varphi_n)$ is a morphism of n -angles.
- (2) The n - Σ -sequence

$$\begin{array}{ccccccc}
 A_3 \xrightarrow{\begin{bmatrix} \alpha_3 \\ \varphi_3 \end{bmatrix}} A_4 \oplus B_3 \xrightarrow{\begin{bmatrix} -\alpha_4 & 0 \\ \varphi_4 & -\beta_3 \\ \psi_2 & \psi_1 \end{bmatrix}} A_5 \oplus B_4 \oplus C_3 \xrightarrow{\begin{bmatrix} -\alpha_5 & 0 & 0 \\ -\varphi_5 & -\beta_4 & 0 \\ \psi_4 & \psi_3 & \gamma_3 \end{bmatrix}} A_6 \oplus B_5 \oplus C_4 \xrightarrow{\begin{bmatrix} -\alpha_6 & 0 & 0 \\ \varphi_6 & -\beta_5 & 0 \\ \psi_6 & \psi_5 & \gamma_4 \end{bmatrix}} \cdots \\
 \xrightarrow{\begin{bmatrix} -\alpha_{n-1} & 0 & 0 \\ (-1)^{n+1}\varphi_{n-1} & -\beta_{n-2} & 0 \\ \psi_{2n-8} & \psi_{2n-9} & \gamma_{n-3} \end{bmatrix}} A_n \oplus B_{n-1} \oplus C_{n-2} \xrightarrow{\begin{bmatrix} (-1)^n\varphi_n & -\beta_{n-1} & 0 \\ \psi_{2n-6} & \psi_{2n-7} & \gamma_{n-2} \end{bmatrix}} B_n \oplus C_{n-1} \xrightarrow{[\psi_{2n-5} \ \gamma_{n-1}]} C_n \xrightarrow{\Sigma\alpha_2 \circ \gamma_n} \Sigma A_3
 \end{array}$$

is an n -angle, and $\gamma_n \circ \psi_{2n-5} = \Sigma\alpha_1 \circ \beta_n$.

For small values of n , objects A_i, B_i, C_i with $i > n$ appearing in the axiom should be interpreted as zero objects (and so should objects C_i with $i < 3$). Specifically, when $n = 3$, that is, when \mathcal{C} is a triangulated category, the triangle in (2) becomes

$$A_3 \xrightarrow{\varphi_3} B_3 \xrightarrow{\psi_1} C_3 \xrightarrow{\Sigma\alpha_2 \circ \gamma_3} \Sigma A_3$$

and for $n = 4$, the 4-angle in (2) becomes

$$A_3 \xrightarrow{\begin{bmatrix} \alpha_3 \\ \varphi_3 \end{bmatrix}} A_4 \oplus B_3 \xrightarrow{\begin{bmatrix} \varphi_4 & -\beta_3 \\ \psi_2 & \psi_1 \end{bmatrix}} B_4 \oplus C_3 \xrightarrow{[\psi_3 \ \gamma_3]} C_4 \xrightarrow{\Sigma\alpha_2 \circ \gamma_4} \Sigma A_3.$$

Our aim is to prove that axiom (N4) may be replaced by the new axiom (N4*). In other words, we shall prove that if our category \mathcal{C} is pre-triangulated (that is, \mathcal{C} satisfies (N1), (N2) and (N3)), then it satisfies (N4) if and only if it satisfies (N4*). In order to prove this, we need the following lemma.

Lemma 4.1. *Suppose \mathcal{C} is n -angulated, and let*

$$\begin{array}{ccccccccccc}
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
 \parallel & & \downarrow \varphi_2 & & & & & & & & \parallel \\
 A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma A_1
 \end{array}$$

be a commutative diagram whose rows are n -angles. Apply axiom (N4) and complete the diagram to a morphism

$$\begin{array}{ccccccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
\parallel & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \parallel \\
A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma A_1
\end{array}$$

of n -angles, in such a way that the mapping cone is also an n -angle. Then the n - Σ -sequence

$$\begin{aligned}
A_2 & \xrightarrow{\begin{bmatrix} -\alpha_2 \\ \varphi_2 \end{bmatrix}} A_3 \oplus B_2 \xrightarrow{\begin{bmatrix} \alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} A_4 \oplus B_3 \xrightarrow{\begin{bmatrix} \alpha_4 & 0 \\ -\varphi_4 & \beta_3 \end{bmatrix}} \cdots \\
& \cdots \xrightarrow{\begin{bmatrix} \alpha_{n-1} & 0 \\ (-1)^n \varphi_{n-1} & \beta_{n-2} \end{bmatrix}} A_n \oplus B_{n-1} \xrightarrow{\begin{bmatrix} (-1)^{n+1} \varphi_n & \beta_{n-1} \end{bmatrix}} B_n \xrightarrow{\Sigma \alpha_1 \circ \beta_n} \Sigma A_2
\end{aligned}$$

is an n -angle.

Proof. The mapping cone is the middle n - Σ -sequence in the direct sum diagram

$$\begin{array}{ccccccccccc}
A_2 & \xrightarrow{\begin{bmatrix} -\alpha_2 \\ \varphi_2 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{\begin{bmatrix} \alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} & \cdots & \xrightarrow{\begin{bmatrix} \alpha_{n-1} & 0 \\ (-1)^n \varphi_{n-1} & \beta_{n-2} \end{bmatrix}} & A_n \oplus B_{n-1} & \xrightarrow{\begin{bmatrix} (-1)^{n+1} \varphi_n & \beta_{n-1} \end{bmatrix}} & B_n & \xrightarrow{\Sigma \alpha_1 \circ \beta_n} & \Sigma A_2 \\
\downarrow \begin{bmatrix} 1 \\ 0 \end{bmatrix} & & \downarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} & & & & \downarrow \begin{bmatrix} (-1)^{n+1} & 0 \\ 0 & 1 \end{bmatrix} & & \downarrow \begin{bmatrix} -\beta_n \\ 1 \end{bmatrix} & & \downarrow \begin{bmatrix} 1 \\ 0 \end{bmatrix} \\
A_2 \oplus A_1 & \xrightarrow{\begin{bmatrix} -\alpha_2 & 0 \\ \varphi_2 & \beta_1 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{\begin{bmatrix} -\alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} & \cdots & \xrightarrow{\begin{bmatrix} -\alpha_{n-1} & 0 \\ \varphi_{n-1} & \beta_{n-2} \end{bmatrix}} & A_n \oplus B_{n-1} & \xrightarrow{\begin{bmatrix} -\alpha_n & 0 \\ \varphi_n & \beta_{n-1} \end{bmatrix}} & \Sigma A_1 \oplus B_n & \xrightarrow{\begin{bmatrix} -\Sigma \alpha_1 & 0 \\ 1 & \beta_n \end{bmatrix}} & \Sigma A_2 \oplus \Sigma A_1 \\
\downarrow \begin{bmatrix} 1 & \alpha_1 \end{bmatrix} & & \downarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} & & & & \downarrow \begin{bmatrix} (-1)^{n+1} & 0 \\ 0 & 1 \end{bmatrix} & & \downarrow \begin{bmatrix} 0 & 1 \end{bmatrix} & & \downarrow \begin{bmatrix} 1 & \Sigma \alpha_1 \end{bmatrix} \\
A_2 & \xrightarrow{\begin{bmatrix} -\alpha_2 \\ \varphi_2 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{\begin{bmatrix} \alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} & \cdots & \xrightarrow{\begin{bmatrix} \alpha_{n-1} & 0 \\ (-1)^n \varphi_{n-1} & \beta_{n-2} \end{bmatrix}} & A_n \oplus B_{n-1} & \xrightarrow{\begin{bmatrix} (-1)^{n+1} \varphi_n & \beta_{n-1} \end{bmatrix}} & B_n & \xrightarrow{\Sigma \alpha_1 \circ \beta_n} & \Sigma A_2
\end{array}$$

Therefore, by axiom (N1)(a), the top (bottom) row is also an n -angle. \square

Now we prove that axioms (N4) and (N4*) are equivalent. We do this in two steps, showing first that axiom (N4) implies axiom (N4*).

Theorem 4.2. *If \mathcal{N} is a collection of n - Σ -sequences in \mathcal{C} satisfying axioms (N1), (N2), (N3) and (N4), then it also satisfies (N4*).*

Proof. Suppose we are given a commutative diagram

$$\begin{array}{ccccccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-2}} & A_{n-1} & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
\parallel & & \downarrow \varphi_2 & & & & & & & & & & \parallel \\
A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-2}} & B_{n-1} & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma A_1
\end{array}$$

where the two rows are n -angles, and where the map φ_1 is an isomorphism. Furthermore, let

$$A_2 \xrightarrow{\varphi_2} B_2 \xrightarrow{\gamma_2} C_3 \xrightarrow{\gamma_3} \cdots \xrightarrow{\gamma_{n-1}} C_n \xrightarrow{\gamma_n} \Sigma A_2$$

be an n -angle. Apply axiom (N4) and complete the given diagram to a morphism $(1, \varphi_2, \varphi_3, \dots, \varphi_n)$ of n -angles, in such a way that the mapping cone is an n -angle. Then the first part of axiom (N4*) is already satisfied.

By Lemma 4.1, the n - Σ -sequence

$$A_2 \xrightarrow{\begin{bmatrix} -\alpha_2 \\ \varphi_2 \end{bmatrix}} A_3 \oplus B_2 \xrightarrow{\begin{bmatrix} \alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} A_4 \oplus B_3 \xrightarrow{\begin{bmatrix} \alpha_4 & 0 \\ -\varphi_4 & \beta_3 \end{bmatrix}} \cdots \xrightarrow{\begin{bmatrix} \alpha_{n-1} & 0 \\ (-1)^n \varphi_{n-1} & \beta_{n-2} \end{bmatrix}} A_n \oplus B_{n-1} \xrightarrow{\begin{bmatrix} (-1)^{n+1} \varphi_n & \beta_{n-1} \end{bmatrix}} B_n \xrightarrow{\Sigma \alpha_1 \circ \beta_n} \Sigma A_2$$

is an n -angle. Then by axiom (N4) again, there exist morphisms $\psi_3, \psi_4, \dots, \psi_{2n-5}$ such that the mapping cone of the morphism

$$\begin{array}{ccccccccccccccc}
 A_2 & \xrightarrow{\begin{bmatrix} -\alpha_2 \\ \varphi_2 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{\begin{bmatrix} \alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} & A_4 \oplus B_3 & \xrightarrow{\begin{bmatrix} \alpha_4 & 0 \\ -\varphi_4 & \beta_3 \end{bmatrix}} & \cdots & \xrightarrow{\begin{bmatrix} \alpha_{n-1} & 0 \\ (-1)^n \varphi_{n-1} & \beta_{n-2} \end{bmatrix}} & A_n \oplus B_{n-1} & \xrightarrow{\begin{bmatrix} (-1)^{n+1} \varphi_n & \beta_{n-1} \end{bmatrix}} & B_n & \xrightarrow{\Sigma \alpha_1 \circ \beta_n} & \Sigma A_2 \\
 \parallel & & \downarrow [0 \ 1] & & \downarrow [\psi_2 \ \psi_1] & & & & \downarrow [\psi_{2n-6} \ \psi_{2n-7}] & & \downarrow \psi_{2n-5} & & \parallel \\
 A_2 & \xrightarrow{\varphi_2} & B_2 & \xrightarrow{\gamma_2} & C_3 & \xrightarrow{\gamma_3} & \cdots & \xrightarrow{\gamma_{n-2}} & C_{n-1} & \xrightarrow{\gamma_{n-1}} & C_n & \xrightarrow{\gamma_n} & \Sigma A_2
 \end{array}$$

is an n -angle. In other words, the n - Σ -sequence

$$\begin{array}{c}
 A_3 \oplus B_2 \oplus A_2 \xrightarrow{\begin{bmatrix} -\alpha_3 & 0 & 0 \\ -\varphi_3 & -\beta_2 & 0 \\ 0 & 1 & \varphi_2 \end{bmatrix}} A_4 \oplus B_3 \oplus B_2 \xrightarrow{\mu_1} A_5 \oplus B_4 \oplus C_3 \xrightarrow{\mu_2} \cdots \\
 \xrightarrow{\mu_{n-4}} A_n \oplus B_{n-1} \oplus C_{n-2} \xrightarrow{\begin{bmatrix} (-1)^n \varphi_n & -\beta_{n-1} & 0 \\ \psi_{2n-6} & \psi_{2n-7} & \gamma_{n-2} \end{bmatrix}} B_n \oplus C_{n-1} \xrightarrow{\begin{bmatrix} -\Sigma \alpha_1 \circ \beta_n & 0 \\ \psi_{2n-5} & \gamma_{n-1} \end{bmatrix}} \Sigma A_2 \oplus C_n \xrightarrow{\begin{bmatrix} \Sigma \alpha_2 & 0 \\ -\Sigma \varphi_2 & 0 \\ 1 & \gamma_n \end{bmatrix}} \Sigma A_3 \oplus \Sigma B_2 \oplus \Sigma A_2
 \end{array}$$

is an n -angle, where μ_i is the matrix

$$\mu_i = \begin{bmatrix} -\alpha_{i+3} & 0 & 0 \\ (-1)^{i+1} \varphi_{i+3} & -\beta_{i+2} & 0 \\ \psi_{2i} & \psi_{2i-1} & \gamma_{i+1} \end{bmatrix}.$$

This n -angle is the middle n - Σ -sequence in the direct sum diagram

$$\begin{array}{ccccccccccc}
 A_3 & \xrightarrow{\begin{bmatrix} \alpha_3 \\ \varphi_3 \end{bmatrix}} & A_4 \oplus B_3 & \xrightarrow{\begin{bmatrix} -\alpha_4 & 0 \\ \varphi_4 & -\beta_3 \\ \psi_2 & \psi_1 \end{bmatrix}} & A_5 \oplus B_4 \oplus C_3 & \xrightarrow{\mu_2} & A_6 \oplus B_5 \oplus C_4 & \xrightarrow{\mu_3} & \cdots \\
 \downarrow \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} & & \downarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} & & \parallel & & \parallel & & \\
 A_3 \oplus B_2 \oplus A_2 & \xrightarrow{\begin{bmatrix} -\alpha_3 & 0 & 0 \\ -\varphi_3 & -\beta_2 & 0 \\ 0 & 1 & \varphi_2 \end{bmatrix}} & A_4 \oplus B_3 \oplus B_2 & \xrightarrow{\mu_1} & A_5 \oplus B_4 \oplus C_3 & \xrightarrow{\mu_2} & A_6 \oplus B_5 \oplus C_4 & \xrightarrow{\mu_3} & \cdots \\
 \downarrow [-1 \ 0 \ \alpha_2] & & \downarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & \beta_2 \end{bmatrix} & & \parallel & & \parallel & & \\
 A_3 & \xrightarrow{\begin{bmatrix} \alpha_3 \\ \varphi_3 \end{bmatrix}} & A_4 \oplus B_3 & \xrightarrow{\begin{bmatrix} -\alpha_4 & 0 \\ \varphi_4 & -\beta_3 \\ \psi_2 & \psi_1 \end{bmatrix}} & A_5 \oplus B_4 \oplus C_3 & \xrightarrow{\mu_2} & A_6 \oplus B_5 \oplus C_4 & \xrightarrow{\mu_3} & \cdots
 \end{array}$$

$$\begin{array}{ccccccc}
 \xrightarrow{\mu_{n-4}} A_n \oplus B_{n-1} \oplus C_{n-2} \xrightarrow{\begin{bmatrix} (-1)^n \varphi_n & -\beta_{n-1} & 0 \\ \psi_{2n-6} & \psi_{2n-7} & \gamma_{n-2} \end{bmatrix}} B_n \oplus C_{n-1} \xrightarrow{[\psi_{2n-5} \ \gamma_{n-1}]} C_n \xrightarrow{\Sigma \alpha_2 \circ \gamma_n} \Sigma A_3 \\
 \parallel & & \parallel & & \downarrow \begin{bmatrix} -\gamma_n \\ 1 \end{bmatrix} & & \downarrow \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix} \\
 \xrightarrow{\mu_{n-4}} A_n \oplus B_{n-1} \oplus C_{n-2} \xrightarrow{\begin{bmatrix} (-1)^n \varphi_n & -\beta_{n-1} & 0 \\ \psi_{2n-6} & \psi_{2n-7} & \gamma_{n-2} \end{bmatrix}} B_n \oplus C_{n-1} \xrightarrow{\begin{bmatrix} -\Sigma \alpha_1 \circ \beta_n & 0 \\ \psi_{2n-5} & \gamma_{n-1} \end{bmatrix}} \Sigma A_2 \oplus C_n \xrightarrow{\begin{bmatrix} \Sigma \alpha_2 & 0 \\ -\Sigma \varphi_2 & 0 \\ 1 & \gamma_n \end{bmatrix}} \Sigma A_3 \oplus \Sigma B_2 \oplus \Sigma A_2 \\
 \parallel & & \parallel & & \downarrow [0 \ 1] & & \downarrow [-1 \ 0 \ \Sigma \alpha_2] \\
 \xrightarrow{\mu_{n-4}} A_n \oplus B_{n-1} \oplus C_{n-2} \xrightarrow{\begin{bmatrix} (-1)^n \varphi_n & -\beta_{n-1} & 0 \\ \psi_{2n-6} & \psi_{2n-7} & \gamma_{n-2} \end{bmatrix}} B_n \oplus C_{n-1} \xrightarrow{[\psi_{2n-5} \ \gamma_{n-1}]} C_n \xrightarrow{\Sigma \alpha_2 \circ \gamma_n} \Sigma A_3
 \end{array}$$

Consequently, by (N1)(a), the top (bottom) n - Σ -sequence is an n -angle. Moreover, the commutativity of the square Ω implies that $\gamma_n \circ \psi_{2n-5} = \Sigma \alpha_1 \circ \beta_n$. This shows that the second part of axiom (N4*) is satisfied. \square

We now prove the converse to Theorem 4.2, namely that the octahedral axiom (N4*) implies axiom (N4).

Theorem 4.3. *If \mathcal{N} is a collection of n - Σ -sequences in \mathcal{C} satisfying axioms (N1), (N2), (N3) and (N4*), then it also satisfies (N4).*

Proof. Given a commutative diagram

$$\begin{array}{ccccccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
\downarrow \varphi_1 & & \downarrow \varphi_2 & & & & & & & & \downarrow \Sigma \varphi_1 \\
B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1
\end{array}$$

where the two rows are n -angles: we denote these by A_\bullet and B_\bullet . We want to prove that we can complete the above diagram to a morphism of n -angles in such a way that the mapping cone of that morphism is again an n -angle.

From the given diagram we build the diagram

$$\begin{array}{ccccccccccc}
A_1 \oplus B_1 & \xrightarrow{\begin{bmatrix} 0 & 0 \\ (-1)^n \alpha_1 & 0 \\ 0 & -1 \end{bmatrix}} & B_2 \oplus A_2 \oplus B_1 & \xrightarrow{\begin{bmatrix} 0 & -\alpha_2 & 0 \\ 1 & 0 & 0 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{[-\alpha_3 \ 0]} & A_4 & \xrightarrow{\alpha_4} & \cdots \\
\parallel & & \downarrow [1 \ -\varphi_2 \ -\beta_1] & & & & & & & & \\
A_1 \oplus B_1 & \xrightarrow{[(-1)^{n+1} \varphi_2 \circ \alpha_1 \ \beta_1]} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{-\beta_3} & B_4 & \xrightarrow{-\beta_4} & \cdots \\
& & \downarrow & & & & & & & & \\
& & 0 & \xrightarrow{\alpha_{n-2}} & A_{n-1} & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\begin{bmatrix} (-1)^n \alpha_n \\ 0 \end{bmatrix}} & \Sigma A_1 \oplus \Sigma B_1 & & \\
& & \downarrow & & & & & & & & \\
& & \vdots & & \cdots & \xrightarrow{-\beta_{n-2}} & B_{n-1} & \xrightarrow{\begin{bmatrix} 0 \\ -\beta_{n-1} \end{bmatrix}} & \Sigma A_1 \oplus B_n & \xrightarrow{\begin{bmatrix} (-1)^{-1} \Sigma \varphi_1 & 0 \\ (-1)^{n+1} \beta_n \end{bmatrix}} & \Sigma A_1 \oplus \Sigma B_1 \\
& & \downarrow & & & & & & & & \\
& & 0 & & & & & & & & \\
& & \downarrow & & & & & & & & \\
& & \Sigma A_2 \oplus \Sigma B_1 & & & & & & & & \\
& & \downarrow \begin{bmatrix} (-1)^n \Sigma \varphi_2 & (-1)^n \Sigma \beta_1 \\ (-1)^n & 0 \\ 0 & (-1)^n \end{bmatrix} & & & & & & & & \\
& & \Sigma B_2 \oplus \Sigma A_2 \oplus \Sigma B_1 & & & & & & & &
\end{array}$$

in which the top left square commutes. Let X_\bullet , Y_\bullet and Z_\bullet denote the three n - Σ -sequences

$$\begin{aligned}
X_\bullet &: B_2 \oplus A_2 \oplus B_1 \xrightarrow{[1 \ -\varphi_2 \ -\beta_1]} B_2 \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow \Sigma A_2 \oplus \Sigma B_1 \xrightarrow{\begin{bmatrix} (-1)^n \Sigma \varphi_2 & (-1)^n \Sigma \beta_1 \\ (-1)^n & 0 \\ 0 & (-1)^n \end{bmatrix}} \Sigma B_2 \oplus \Sigma A_2 \oplus \Sigma B_1, \\
Y_\bullet &: A_1 \oplus B_1 \xrightarrow{\begin{bmatrix} 0 & 0 \\ (-1)^n \alpha_1 & 0 \\ 0 & -1 \end{bmatrix}} B_2 \oplus A_2 \oplus B_1 \xrightarrow{\begin{bmatrix} 0 & -\alpha_2 & 0 \\ 1 & 0 & 0 \end{bmatrix}} A_3 \oplus B_2 \xrightarrow{[-\alpha_3 \ 0]} A_4 \xrightarrow{\alpha_4} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\begin{bmatrix} (-1)^n \alpha_n \\ 0 \end{bmatrix}} \Sigma A_1 \oplus \Sigma B_1, \\
Z_\bullet &: A_1 \oplus B_1 \xrightarrow{[(-1)^{n+1} \varphi_2 \circ \alpha_1 \ \beta_1]} B_2 \xrightarrow{\beta_2} B_3 \xrightarrow{-\beta_3} \cdots \xrightarrow{-\beta_{n-2}} B_{n-1} \xrightarrow{\begin{bmatrix} 0 \\ -\beta_{n-1} \end{bmatrix}} \Sigma A_1 \oplus B_n \xrightarrow{\begin{bmatrix} -1 & 0 \\ (-1)^{n+1} \Sigma \varphi_1 & (-1)^{n+1} \beta_n \end{bmatrix}} \Sigma A_1 \oplus \Sigma B_1,
\end{aligned}$$

respectively. In order to apply (N4*) we need to prove that these n - Σ -sequences are n -angles.

It can easily be shown that X_\bullet is isomorphic to the direct sum of the trivial n -angle on B_2 and the left rotations of the trivial n -angles on A_2 and B_1 . Next, the n - Σ -sequence Y_\bullet is isomorphic to the direct sum of the n -angle A_\bullet , the trivial n -angle on B_1 and the right rotation of the trivial n -angle on B_2 . Similarly, the n - Σ -sequence Z_\bullet is isomorphic to the direct sum of the n -angle B_\bullet and the left rotation of the trivial n -angle on A_1 . Hence, by (N1)(a) it follows that X_\bullet , Y_\bullet and Z_\bullet are n -angles.

Since X_\bullet , Y_\bullet and Z_\bullet are n -angles, we may apply axiom (N4*) to the above diagram. Consequently, there exist morphisms $\sigma_3, \sigma_4, \dots, \sigma_n$ and a morphism ψ_{2n-5} with the following three properties:

- (1) the sequence $(1, [1 \ -\varphi_2 \ -\beta_1], \sigma_3, \sigma_4, \dots, \sigma_n)$ is a morphism $Y_\bullet \rightarrow Z_\bullet$ of n -angles,
- (2) ψ_{2n-5} is a morphism $\Sigma A_1 \oplus B_n \rightarrow \Sigma A_2 \oplus \Sigma B_1$ with

$$\begin{bmatrix} (-1)^n \Sigma \varphi_2 & (-1)^n \Sigma \beta_1 \\ (-1)^n & 0 \\ 0 & (-1)^n \end{bmatrix} \circ \psi_{2n-5} = \begin{bmatrix} 0 & 0 \\ (-1)^n \Sigma \alpha_1 & 0 \\ 0 & -1 \end{bmatrix} \circ \begin{bmatrix} -1 & 0 \\ (-1)^{n+1} \Sigma \varphi_1 & (-1)^{n+1} \beta_n \end{bmatrix}.$$

(3) the n - Σ -sequence

$$A_3 \oplus B_2 \xrightarrow{\begin{bmatrix} -\alpha_3 & 0 \\ \sigma_{3,1} & \sigma_{3,2} \end{bmatrix}} A_4 \oplus B_3 \xrightarrow{\begin{bmatrix} -\alpha_4 & 0 \\ \sigma_4 & \beta_3 \end{bmatrix}} \dots \xrightarrow{\psi_{2n-5}} \Sigma A_2 \oplus \Sigma B_1 \xrightarrow{\begin{bmatrix} (-1)^{n+1}\Sigma\alpha_2 & 0 \\ (-1)^n\Sigma\varphi_2 & (-1)^n\Sigma\beta_1 \end{bmatrix}} \Sigma A_3 \oplus \Sigma B_2$$

is an n -angle.

Observe that the n -angle X_\bullet consists of the zero object at positions 3 through $n-1$, thus the other morphisms $\psi_1, \psi_2, \dots, \psi_{2n-6}$ given by (N4*) are all zero.

From property (1) the diagram

$$\begin{array}{ccccccc} A_1 \oplus B_1 & \xrightarrow{\begin{bmatrix} 0 & 0 \\ (-1)^n\alpha_1 & 0 \\ 0 & -1 \end{bmatrix}} & B_2 \oplus A_2 \oplus B_1 & \xrightarrow{\begin{bmatrix} 0 & -\alpha_2 & 0 \\ 1 & 0 & 0 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{[-\alpha_3 \ 0]} & A_4 & \xrightarrow{\alpha_4} & \dots \\ \parallel & & \downarrow [1 \ -\varphi_2 \ -\beta_1] & & \downarrow \sigma_3 & & \downarrow \sigma_4 & & \\ A_1 \oplus B_1 & \xrightarrow{[(-1)^{n+1}\varphi_2 \circ \alpha_1 \ \beta_1]} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{-\beta_3} & B_4 & \xrightarrow{-\beta_4} & \dots \end{array}$$

$$\begin{array}{ccccccc} & & \xrightarrow{\alpha_{n-2}} & A_{n-1} & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\begin{bmatrix} (-1)^n\alpha_n \\ 0 \end{bmatrix}} & \Sigma A_1 \oplus \Sigma B_1 \\ & & & \downarrow \sigma_{n-1} & & \downarrow \sigma_n & & \parallel \\ & & \xrightarrow{-\beta_{n-2}} & B_{n-1} & \xrightarrow{\begin{bmatrix} 0 \\ -\beta_{n-1} \end{bmatrix}} & \Sigma A_1 \oplus B_n & \xrightarrow{\begin{bmatrix} (-1)^{n+1}\Sigma\varphi_1 & 0 \\ (-1)^{n+1}\beta_n \end{bmatrix}} & \Sigma A_1 \oplus \Sigma B_1 \end{array}$$

is commutative. Using the commutativity, we can conclude that

$$\begin{aligned} \sigma_3 &= [\varphi_3 \ \beta_2], \\ \sigma_4 &= \varphi_4, \\ \sigma_5 &= -\varphi_5 \\ &\vdots \\ \sigma_{n-1} &= (-1)^{n-1}\varphi_{n-1}, \\ \sigma_n &= \begin{bmatrix} (-1)^{n+1}\alpha_n \\ (-1)^n\varphi_n \end{bmatrix} \end{aligned}$$

for some morphisms $A_i \xrightarrow{\varphi_i} B_i$ ($3 \leq i \leq n$) making the sequence $\varphi = (\varphi_1, \varphi_2, \dots, \varphi_n)$ into a morphism $A_\bullet \xrightarrow{\varphi} B_\bullet$ of n -angles.

Next, consider the morphism ψ_{2n-5} . Using property (2), we see that

$$\begin{aligned} \begin{bmatrix} (-1)^n\Sigma\varphi_2 & (-1)^n\Sigma\beta_1 \\ (-1)^n & 0 \\ 0 & (-1)^n \end{bmatrix} \circ \psi_{2n-5} &= \begin{bmatrix} 0 & 0 \\ (-1)^n\Sigma\alpha_1 & 0 \\ 0 & -1 \end{bmatrix} \circ \begin{bmatrix} -1 & 0 \\ (-1)^{n+1}\Sigma\varphi_1 & (-1)^{n+1}\beta_n \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ (-1)^{n+1}\Sigma\alpha_1 & 0 \\ (-1)^n\Sigma\varphi_1 & (-1)^n\beta_n \end{bmatrix}. \end{aligned}$$

Thus the morphism ψ_{2n-5} is given by the matrix

$$\psi_{2n-5} = \begin{bmatrix} -\Sigma\alpha_1 & 0 \\ \Sigma\varphi_1 & \beta_n \end{bmatrix}.$$

Finally, from property (3) and what we have shown so far, the n - Σ -sequence

$$A_3 \oplus B_2 \xrightarrow{\begin{bmatrix} -\alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} A_4 \oplus B_3 \xrightarrow{\begin{bmatrix} -\alpha_4 & 0 \\ \varphi_4 & \beta_3 \end{bmatrix}} \dots \xrightarrow{\begin{bmatrix} -\Sigma\alpha_1 & 0 \\ \Sigma\varphi_1 & \beta_n \end{bmatrix}} \Sigma A_2 \oplus \Sigma B_1 \xrightarrow{\begin{bmatrix} (-1)^{n+1}\Sigma\alpha_2 & 0 \\ (-1)^n\Sigma\varphi_2 & (-1)^n\Sigma\beta_1 \end{bmatrix}} \Sigma A_3 \oplus \Sigma B_2$$

is an n -angle. Its right rotation

$$A_2 \oplus B_1 \xrightarrow{\begin{bmatrix} -\alpha_2 & 0 \\ \varphi_2 & \beta_1 \end{bmatrix}} A_3 \oplus B_2 \xrightarrow{\begin{bmatrix} -\alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{bmatrix}} \dots \xrightarrow{\begin{bmatrix} -\alpha_n & 0 \\ \varphi_n & \beta_{n-1} \end{bmatrix}} \Sigma A_1 \oplus B_n \xrightarrow{\begin{bmatrix} -\Sigma\alpha_1 & 0 \\ \Sigma\varphi_1 & \beta_n \end{bmatrix}} \Sigma A_2 \oplus \Sigma B_1$$

is the mapping cone of φ , and this is an n -angle by axiom (N2). This completes the proof. \square

Collecting Theorem 4.2 and Theorem 4.3 gives the following.

Theorem 4.4. *If \mathcal{N} is a collection of n - Σ -sequences satisfying axioms (N1), (N2) and (N3), then the following are equivalent:*

- (1) \mathcal{N} satisfies (N4),
- (2) \mathcal{N} satisfies (N4*).

We now discuss the case when $n = 3$, that is, when our category \mathcal{C} is a triangulated category. In this case, the classical octahedral axiom, which was introduced by Verdier in [9, 10], is the following:

(TR4) Given a commutative diagram

$$\begin{array}{ccccccc}
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \Sigma A_1 \\
 \parallel & & \downarrow \varphi_2 & & & & \parallel \\
 A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \Sigma A_1 \\
 & & \downarrow \gamma_2 & & & & \\
 & & C_3 & & & & \\
 & & \downarrow \gamma_3 & & & & \\
 & & \Sigma A_2 & & & &
 \end{array}$$

in which the top rows and second column are triangles. Then there exist morphisms $A_3 \xrightarrow{\varphi_3} B_3$ and $B_3 \xrightarrow{\psi_1} C_3$ with the following properties: the diagram

$$\begin{array}{ccccccc}
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \Sigma A_1 \\
 \parallel & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \parallel \\
 A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \Sigma A_1 \\
 & & \downarrow \gamma_2 & \Theta & \downarrow \psi_1 & & \\
 & & C_3 & \xlongequal{\quad} & C_3 & & \\
 & & \downarrow \gamma_3 & & \downarrow \Sigma\alpha_2 \circ \gamma_3 & & \\
 & & \Sigma A_2 & \xrightarrow{\Sigma\alpha_2} & \Sigma A_3 & &
 \end{array}$$

is commutative, the third column is a triangle, and $\gamma_3 \circ \psi_1 = \Sigma\alpha_1 \circ \beta_3$.

This is almost the same as our axiom (N4*): there is one difference. Namely, axiom (N4*) does *not* guarantee that the square Θ commutes. However, when $n = 3$ and we start with the diagram given in (TR4), then in the proof of Theorem 4.2 we obtain the commutative diagram

$$\begin{array}{ccccccc}
 A_2 & \xrightarrow{\begin{bmatrix} -\alpha_2 \\ \varphi_2 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{[\varphi_3 \ \beta_2]} & B_3 & \xrightarrow{\Sigma\alpha_1 \circ \beta_3} & \Sigma A_2 \\
 \parallel & & \downarrow [0 \ 1] & & \downarrow \psi_1 & & \parallel \\
 A_2 & \xrightarrow{\varphi_2} & B_2 & \xrightarrow{\gamma_2} & C_3 & \xrightarrow{\gamma_3} & \Sigma A_2
 \end{array}$$

The commutativity of the middle square implies that the square Θ in (TR4) commutes. Therefore, we recover the original octahedral axiom (TR4) from axioms (N1), (N2), (N3) and (N4). Conversely, Neeman proves in [6, Theorem 1.8] that axioms (N1), (N2), (N3) and (TR4) together imply axiom (N4). Consequently, when $n = 3$ and the collection \mathcal{N} of 3- Σ -sequences satisfies axioms (N1), (N2) and (N3), then the following are equivalent:

- (1) \mathcal{N} satisfies (N4),
- (2) \mathcal{N} satisfies (TR4),
- (3) \mathcal{N} satisfies (N4*).

We end this section with a discussion of homotopy cartesian diagrams. Recall that when $n = 3$, then a commutative square

$$\begin{array}{ccc}
 A_1 & \xrightarrow{\alpha} & A_2 \\
 \downarrow \varphi_1 & & \downarrow \varphi_2 \\
 B_1 & \xrightarrow{\beta} & B_2
 \end{array}$$

is *homotopy cartesian* if there exists a triangle

$$A_1 \xrightarrow{\begin{bmatrix} -\alpha \\ \varphi_1 \end{bmatrix}} A_2 \oplus B_1 \xrightarrow{[\varphi_2 \ \beta]} B_2 \xrightarrow{\partial} \Sigma A_1$$

for some morphism $B_2 \xrightarrow{\partial} \Sigma A_1$. Now let (TR4*) be the axiom which is the same as (TR4), but with the additional requirement that the commutative square

$$\begin{array}{ccc} A_2 & \xrightarrow{\alpha_2} & A_3 \\ \downarrow \varphi_2 & & \downarrow \varphi_3 \\ B_2 & \xrightarrow{\beta_2} & B_3 \end{array}$$

is homotopy cartesian. Neeman shows in [6, 7] that (TR4) is equivalent to the stronger (TR4*). Consequently, the axioms (N4), (N4*), (TR4) and (TR4*) are all equivalent.

Now let \mathcal{C} be n -angulated. Motivated by the above, we say that a commutative diagram

$$\begin{array}{ccccccccc} A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & \cdots & \xrightarrow{\alpha_{n-3}} & A_{n-2} & \xrightarrow{\alpha_{n-2}} & A_{n-1} \\ \downarrow \varphi_1 & & \downarrow \varphi_2 & & & & \downarrow \varphi_{n-2} & & \downarrow \varphi_{n-1} \\ B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & \cdots & \xrightarrow{\beta_{n-3}} & B_{n-2} & \xrightarrow{\beta_{n-2}} & B_{n-1} \end{array}$$

is *homotopy cartesian* if the n - Σ -sequence

$$A_1 \xrightarrow{\begin{bmatrix} -\alpha_1 \\ \varphi_1 \end{bmatrix}} A_2 \oplus B_1 \xrightarrow{\begin{bmatrix} \alpha_2 & 0 \\ \varphi_2 & \beta_1 \end{bmatrix}} A_3 \oplus B_2 \xrightarrow{\begin{bmatrix} \alpha_3 & 0 \\ -\varphi_3 & \beta_2 \end{bmatrix}} \cdots \xrightarrow{\begin{bmatrix} \alpha_{n-2} & 0 \\ (-1)^n \varphi_{n-2} & \beta_{n-3} \end{bmatrix}} A_{n-1} \oplus B_{n-2} \xrightarrow{[(-1)^{n+1} \varphi_{n-1} \ \beta_{n-2}]} B_{n-1} \xrightarrow{\partial} \Sigma A_1$$

is an n -angle for some morphism $B_{n-1} \xrightarrow{\partial} \Sigma A_1$. In the proof of Theorem 4.2, when we showed that axiom (N4*) follows from axiom (N4), we proved in addition that the commutative diagram

$$\begin{array}{ccccccccc} A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-2}} & A_{n-1} & \xrightarrow{\alpha_{n-1}} & A_n \\ \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_{n-1} & & \downarrow \varphi_n \\ B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-2}} & B_{n-1} & \xrightarrow{\beta_{n-1}} & B_n \end{array}$$

is homotopy cartesian. In fact, that was precisely Lemma 4.1. Consequently, axiom (N4*) (and then also axiom (N4)) is equivalent to the stronger axiom which requires the above commutative diagram to be homotopy cartesian.

5. AN EXAMPLE: THE GEISS-KELLER-OPPERMANN-CATEGORY

In this section, we recall the example from [2] of an n -angulated category arising from an $(n-2)$ -cluster tilting subcategory of a triangulated category. Let \mathcal{T} be a triangulated category with suspension Σ , and let \mathcal{C} be a full subcategory. Then \mathcal{C} is an $(n-2)$ -cluster tilting subcategory if it satisfies the following two properties:

- (1) \mathcal{C} is functorially finite in \mathcal{T} .
- (2) \mathcal{C} is given by

$$\begin{aligned} \mathcal{C} &= \{X \in \mathcal{T} \mid \text{Hom}_{\mathcal{T}}(X, \Sigma^i C) = 0 \text{ for } 1 \leq i \leq n-3 \text{ and for all } C \in \mathcal{C}\} \\ &= \{X \in \mathcal{T} \mid \text{Hom}_{\mathcal{T}}(\Sigma^i C, X) = 0 \text{ for } 1 \leq i \leq n-3 \text{ and for all } C \in \mathcal{C}\}. \end{aligned}$$

Suppose in addition that \mathcal{C} is closed under the automorphism Σ^{n-2} of \mathcal{T} , and denote Σ^{n-2} by $\widehat{\Sigma}$. Let \mathcal{N} be the collection of n - $\widehat{\Sigma}$ -sequences

$$A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \widehat{\Sigma} A_1$$

in \mathcal{C} such there exists a diagram

$$\begin{array}{ccccccc} & & A_2 & \xrightarrow{\alpha_2} & A_3 & \cdots & A_{n-2} & \xrightarrow{\alpha_{n-2}} & A_{n-1} \\ & \nearrow \alpha_1 & \Delta & \searrow f_1 & \nearrow g_1 & \Delta & \searrow f_2 & \nearrow g_{n-3} & \Delta & \searrow \alpha_{n-1} \\ A_1 & \xleftarrow{\partial_{n-2}} & X_1 & \xleftarrow{\partial_{n-3}} & X_2 & \cdots & X_{n-3} & \xleftarrow{\partial_1} & A_n \end{array}$$

in \mathcal{T} with the following properties:

- (1) Each diagram triangle Δ is a triangle in \mathcal{T} , where a map $X \rightarrow Y$ denotes a map from X to ΣY .
- (2) The other diagram triangles commute.
- (3) The map α_n equals the composition $\Sigma^{n-3}\partial_{n-2} \circ \Sigma^{n-4}\partial_{n-3} \circ \cdots \circ \partial_1$.

Then it is shown in [2, Section 3, Theorem 1] that $(\mathcal{C}, \widehat{\Sigma}, \mathcal{A})$ is an n -angulated category. In what follows, we verify the octahedral axiom (N4*) in the case when $n = 4$. Thus we assume that \mathcal{C} is a 2-cluster tilting subcategory of \mathcal{T} closed under the automorphism Σ^2 , which we denote by $\widehat{\Sigma}$.

Suppose we are given a commutative diagram

$$\begin{array}{ccccccccc}
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & A_4 & \xrightarrow{\alpha_4} & \widehat{\Sigma}A_1 \\
 \parallel & & \downarrow \varphi_2 & & & & & & \parallel \\
 A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & B_4 & \xrightarrow{\beta_4} & \widehat{\Sigma}A_1 \\
 & & \downarrow \gamma_2 & & & & & & \\
 & & C_3 & & & & & & \\
 & & \downarrow \gamma_3 & & & & & & \\
 & & C_4 & & & & & & \\
 & & \downarrow \gamma_4 & & & & & & \\
 & & \widehat{\Sigma}A_2 & & & & & &
 \end{array}$$

in \mathcal{C} whose top rows and second column are 4-angles. We must find morphisms $A_3 \xrightarrow{\varphi_3} B_3, A_4 \xrightarrow{\varphi_4} B_4$ and ψ_1, ψ_2, ψ_3 with the following two properties:

- (1) The sequence $(1, \varphi_2, \varphi_3, \varphi_4)$ is a morphism of 4-angles.
- (2) The 4- $\widehat{\Sigma}$ -sequence

$$A_3 \xrightarrow{\begin{bmatrix} \alpha_3 \\ \varphi_3 \end{bmatrix}} A_4 \oplus B_3 \xrightarrow{\begin{bmatrix} \varphi_4 & -\beta_3 \\ \psi_2 & \psi_1 \end{bmatrix}} B_4 \oplus C_3 \xrightarrow{[\psi_3 \ \gamma_3]} C_4 \xrightarrow{\widehat{\Sigma}\alpha_2 \circ \gamma_4} \widehat{\Sigma}A_3$$

is a 4-angle in \mathcal{C} , and $\gamma_4 \circ \psi_3 = \widehat{\Sigma}\alpha_1 \circ \beta_4$.

We only use the axioms and properties of the underlying triangulated category \mathcal{T} . The three given 4-angles correspond to three diagrams

$$\begin{array}{c}
 \begin{array}{ccccc}
 & & A_2 & \xrightarrow{\alpha_2} & A_3 \\
 & \nearrow \alpha_1 & \Delta_2 & \xrightarrow{f} & \Delta_1 \searrow \alpha_3 \\
 A_1 & & & & X & \xleftarrow{\partial_1} & A_4 \\
 & \xleftarrow{\partial_2} & & & & &
 \end{array} \\
 \\
 \begin{array}{ccccc}
 & & B_2 & \xrightarrow{\beta_2} & B_3 \\
 & \nearrow \beta_1 & \Delta'_2 & \xrightarrow{f'} & \Delta'_1 \searrow \beta_3 \\
 A_1 & & & & Y & \xleftarrow{\partial'_1} & B_4 \\
 & \xleftarrow{\partial'_2} & & & & &
 \end{array} \\
 \\
 \begin{array}{ccccc}
 & & B_2 & \xrightarrow{\gamma_2} & C_3 \\
 & \nearrow \varphi_2 & \Delta''_2 & \xrightarrow{f''} & \Delta''_1 \searrow \gamma_3 \\
 A_2 & & & & Z & \xleftarrow{\partial''_1} & C_4 \\
 & \xleftarrow{\partial''_2} & & & & &
 \end{array}
 \end{array}$$

in \mathcal{T} , that is, each 4-angle is built from two triangles.

Step 1. Choose a morphism $X \xrightarrow{\varphi} Y$ in \mathcal{T} such that the mapping cone of the morphism

$$\begin{array}{ccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{f} & X & \xrightarrow{\partial_2} & \Sigma A_1 \\
\parallel & & \downarrow \varphi_2 & & \downarrow \varphi & & \parallel \\
A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{f'} & Y & \xrightarrow{\partial'_2} & \Sigma A_1
\end{array}$$

is a triangle in \mathcal{T} :

$$A_2 \oplus A_1 \xrightarrow{\begin{bmatrix} -f & 0 \\ \varphi_2 & \beta_1 \end{bmatrix}} X \oplus B_2 \xrightarrow{\begin{bmatrix} -\partial_2 & 0 \\ \varphi & f' \end{bmatrix}} \Sigma A_1 \oplus Y \xrightarrow{\begin{bmatrix} -\Sigma \alpha_1 & 0 \\ 1 & \partial'_2 \end{bmatrix}} \Sigma A_2 \oplus \Sigma A_1.$$

The 3- Σ sequence

$$A_2 \xrightarrow{\begin{bmatrix} -f \\ \varphi_2 \end{bmatrix}} X \oplus B_2 \xrightarrow{[\varphi \ f']} Y \xrightarrow{\Sigma \alpha_1 \circ \partial'_2} \Sigma A_2$$

is a direct summand, and therefore a triangle in \mathcal{T} .

Step 2. By the 3×3 Lemma (cf. [5, Lemma 2.6]), there exists an object W in \mathcal{T} and maps w_1, \dots, w_5 such that the diagram

$$\begin{array}{ccccccc}
& & \Sigma^{-1} A_4 & \xlongequal{\quad} & \Sigma^{-1} A_4 & & \\
& & \downarrow \begin{bmatrix} -\Sigma^{-1} \partial_1 \\ 0 \end{bmatrix} & & \downarrow -\varphi \circ \Sigma^{-1} \partial_1 & & \\
A_2 & \xrightarrow{\begin{bmatrix} -f \\ \varphi_2 \end{bmatrix}} & X \oplus B_2 & \xrightarrow{[\varphi \ f']} & Y & \xrightarrow{\Sigma \alpha_1 \circ \partial'_2} & \Sigma A_2 \\
\parallel & & \downarrow \begin{bmatrix} g & 0 \\ 0 & 1 \end{bmatrix} & & \downarrow w_3 & & \parallel \\
A_2 & \xrightarrow{\begin{bmatrix} -\alpha_2 \\ \varphi_2 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{[w_1 \ w_2]} & W & \xrightarrow{w_4} & \Sigma A_2 \\
& & \downarrow [a_3 \ 0] & & \downarrow w_5 & & \\
& & A_4 & \xlongequal{\quad} & A_4 & &
\end{array}$$

commutes, and all columns and rows are triangles. The second row is the triangle from step 1, whereas the second column is the direct sum of the left rotation of Δ_1 and the trivial triangle on B_2 . By the octahedral axiom, there exist maps w_6, w_7 such that the diagram

$$\begin{array}{ccccccc}
\Sigma^{-1} A_4 & \xrightarrow{-\varphi \circ \Sigma^{-1} \partial_1} & Y & \xrightarrow{w_3} & W & \xrightarrow{w_5} & A_4 \\
\parallel & & \downarrow g' & & \downarrow \begin{bmatrix} w_5 \\ w_6 \end{bmatrix} & & \parallel \\
\Sigma^{-1} A_4 & \xrightarrow{0} & B_3 & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & A_4 \oplus B_3 & \xrightarrow{[1 \ 0]} & A_4 \\
& & \downarrow \beta_3 & & \downarrow [w_7 \ \beta_3] & & \\
& & B_4 & \xlongequal{\quad} & B_4 & & \\
& & \downarrow \partial'_1 & & \downarrow \Sigma w_3 \circ \partial'_1 & & \\
& & \Sigma Y & \xrightarrow{\Sigma w_3} & \Sigma W & &
\end{array}$$

commutes, all rows and columns are triangles, and $\partial'_1 \circ w_7 = -\Sigma \varphi \circ \partial_1$. The first row is a triangle from the diagram above, the second row is a direct sum of trivial triangles, and the second column is Δ'_1 .

Step 3. Define maps

$$\begin{aligned}
A_3 &\xrightarrow{\varphi_3} B_3, & \varphi_3 &:= w_6 \circ w_1 \\
A_4 &\xrightarrow{\varphi_4} B_4, & \varphi_4 &:= -w_7.
\end{aligned}$$

From the diagrams in steps 1 and 2 we obtain the equalities

$$\begin{aligned}
\varphi_3 \circ \alpha_2 &= w_6 \circ w_1 \circ \alpha_2 = w_6 \circ w_2 \circ \varphi_2 = w_6 \circ w_3 \circ f' \circ \varphi_2 = g' \circ f' \circ \varphi_2 = \beta_2 \circ \varphi_2 \\
\varphi_4 \circ \alpha_3 &= \varphi_4 \circ w_5 \circ w_1 = -w_7 \circ w_5 \circ w_1 = \beta_3 \circ w_6 \circ w_1 = \beta_3 \circ \varphi_3 \\
\alpha_4 &= \Sigma \partial_2 \circ \partial_1 = \Sigma \partial'_2 \circ \Sigma \varphi \circ \partial_1 = \Sigma \partial'_2 \circ \partial'_1 \circ (-w_7) = \beta_4 \circ \varphi_4
\end{aligned}$$

of maps in \mathcal{T} . This shows that the diagram

$$\begin{array}{ccccccccc} A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & A_4 & \xrightarrow{\alpha_4} & \widehat{\Sigma}A_1 \\ \parallel & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \downarrow \varphi_3 & & \parallel \\ A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & B_4 & \xrightarrow{\beta_4} & \widehat{\Sigma}A_1 \end{array}$$

is a morphism of 4-angles in \mathcal{C} , hence the first part of axiom (N4*) is satisfied.

Step 4. Choose a morphism $W \xrightarrow{\psi} Z$ in \mathcal{T} such that the mapping cone of the morphism

$$\begin{array}{ccccccc} A_2 & \xrightarrow{\begin{bmatrix} -\alpha_2 \\ \varphi_2 \end{bmatrix}} & A_3 \oplus B_2 & \xrightarrow{\begin{bmatrix} w_1 & w_2 \end{bmatrix}} & W & \xrightarrow{w_4} & \Sigma A_2 \\ \parallel & & \downarrow \begin{bmatrix} 0 & 1 \end{bmatrix} & & \downarrow \psi & & \parallel \\ A_2 & \xrightarrow{\varphi_2} & B_2 & \xrightarrow{f''} & Z & \xrightarrow{\delta_2''} & \Sigma A_2 \end{array}$$

is a triangle in \mathcal{T} :

$$A_3 \oplus B_2 \oplus A_2 \xrightarrow{\begin{bmatrix} -w_1 & -w_2 & 0 \\ 0 & 1 & \varphi_2 \end{bmatrix}} W \oplus B_2 \xrightarrow{\begin{bmatrix} -w_4 & 0 \\ \psi & f'' \end{bmatrix}} \Sigma A_2 \oplus Z \xrightarrow{\begin{bmatrix} \Sigma \alpha_2 & 0 \\ -\Sigma \varphi_2 & 0 \\ 1 & \delta_2'' \end{bmatrix}} \Sigma A_3 \oplus \Sigma B_2 \oplus \Sigma A_2.$$

The top row in the diagram is a triangle from step 2, whereas the bottom row is Δ_2'' . The 3- Σ sequence

$$A_3 \xrightarrow{-w_1} W \xrightarrow{\psi} Z \xrightarrow{-\Sigma \alpha_2 \circ \delta_2''} \Sigma A_3$$

is a direct summand of the mapping cone, and therefore a triangle in \mathcal{T} .

Step 5. By the 3 \times 3 Lemma, there exists an object U in \mathcal{T} and maps u_1, \dots, u_5 such that the diagram

$$\begin{array}{ccccccc} & & \Sigma^{-1}B_4 & \xlongequal{\quad} & \Sigma^{-1}B_4 & & \\ & & \downarrow w_3 \circ \Sigma^{-1}\delta_1' & & \downarrow \psi \circ w_3 \circ \Sigma^{-1}\delta_1' & & \\ A_3 & \xrightarrow{w_1} & W & \xrightarrow{\psi} & Z & \xrightarrow{\Sigma \alpha_2 \circ \delta_2''} & \Sigma A_3 \\ \parallel & & \downarrow \begin{bmatrix} w_5 \\ w_6 \end{bmatrix} & & \downarrow u_3 & & \parallel \\ A_3 & \xrightarrow{\begin{bmatrix} \alpha_3 \\ \varphi_3 \end{bmatrix}} & A_4 \oplus B_3 & \xrightarrow{\begin{bmatrix} u_1 & u_2 \end{bmatrix}} & U & \xrightarrow{u_4} & \Sigma A_3 \\ & & \downarrow \begin{bmatrix} \varphi_4 & -\beta_3 \end{bmatrix} & & \downarrow u_5 & & \\ & & B_4 & \xlongequal{\quad} & B_4 & & \end{array}$$

commutes, and all columns and rows are triangles. The second row is isomorphic to the triangle from step 4, whereas the second column is isomorphic to the right rotation of the third column in the octahedral diagram in step 2. By the octahedral axiom, there exist maps u_6, u_7 such that the diagram

$$\begin{array}{ccccccc} \Sigma^{-1}B_4 & \xrightarrow{\psi \circ w_3 \circ \Sigma^{-1}\delta_1'} & Z & \xrightarrow{u_3} & U & \xrightarrow{u_5} & B_4 \\ \parallel & & \downarrow g'' & & \downarrow \begin{bmatrix} u_5 \\ u_6 \end{bmatrix} & & \parallel \\ \Sigma^{-1}B_4 & \xrightarrow{0} & C_3 & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & B_4 \oplus C_3 & \xrightarrow{\begin{bmatrix} 1 & 0 \end{bmatrix}} & B_4 \\ & & \downarrow \gamma_3 & & \downarrow \begin{bmatrix} u_7 & \gamma_3 \end{bmatrix} & & \\ & & C_4 & \xlongequal{\quad} & C_4 & & \\ & & \downarrow \delta_1'' & & \downarrow \Sigma u_3 \circ \delta_1'' & & \\ & & \Sigma Z & \xrightarrow{\Sigma u_3} & \Sigma U & & \end{array}$$

commutes, all rows and columns are triangles, and $\delta_1'' \circ u_7 = \Sigma \psi \circ \Sigma w_3 \circ \delta_1'$. The first row is a triangle from the diagram above, the second row is a direct sum of trivial triangles, and the second column is Δ_1'' .

