



Norwegian University of  
Science and Technology

Department of Mathematical Sciences

## Examination paper for **ST0103 Statistics with applications**

**Academic contact during examination:** Nikolai Ushakov<sup>a</sup>, Andreas Strand<sup>b</sup>

**Phone:** <sup>a</sup>45128897, <sup>b</sup>90681168

**Examination date:** 16. desember 2017

**Examination time (from–to):** 09:00 – 13:00

**Permitted examination support material:** C: Specific simple calculator. Tabeller og formler i statistikk (Tapir akademisk forlag). One yellow A4 sheet with your own handwritten notes.

**Other information:**

**Language:** English

**Number of pages:** 2

**Number of pages enclosed:** 0

**Checked by:**

Informasjon om trykking av eksamensoppgave	
Originalen er:	
1-sidig <input type="checkbox"/>	2-sidig <input checked="" type="checkbox"/>
sort/hvit <input checked="" type="checkbox"/>	farger <input type="checkbox"/>
skal ha flervalgskjema <input type="checkbox"/>	

---

Date

Signature



**Problem 1** A randomly chosen nest of a bird species contains  $X$  eggs. The probability distribution of  $X$  is as follows:

$x$	1	2	3	4	5	6
$P(X = x)$	0.09	0.38	0.25	0.20	0.06	0.02

Assume that the number of eggs in each nest is independent of each other.

- a) Find the expected value and standard deviation of  $X$ . Find the probability that a randomly chosen nest has 4 or more eggs. Find the conditional probability that a randomly chosen nest has 6 eggs given that it has 4 or more eggs.
- b) We examine 50 randomly chosen nests. Let  $Y$  be the number of these having 4 or more eggs. What is the expected value and standard deviation of  $Y$ ? Find an approximate probability that 10 or more of the 50 nests have at least 4 eggs.
- c) If a nest of this bird species is built in a deciduous tree, the probability is 0.05 that it contains only one egg. If a nest is built in a coniferous tree, the probability is 0.15 that it contains only one egg. The bird species always build the nest in a deciduous tree or in a coniferous tree. What is the probability that a randomly chosen nest is built in a deciduous tree?

**Problem 2**  $Y$  has the normal distribution with the expectation  $\mu$  and variance  $\sigma^2$ .

- a) Find  $P(\mu - 2\sigma < Y < \mu + 2\sigma)$ .
- b) Let  $P(Y < 0) = 0.5$  and  $P(Y < 1) = 0.6915$ . What are  $\mu$  and  $\sigma^2$ ?

**Problem 3** In this exercise we will consider a regression model that is somewhat modified relative to what is discussed in the textbook. Assume that we have pairs of variables

$$(x_1, Y_1), (x_2, Y_2), \dots, (x_n, Y_n)$$

where  $x_1, x_2, \dots, x_n$  are positive and nonstochastic while  $Y_1, Y_2, \dots, Y_n$  are assumed to be independent random variables with

$$Y_i \sim N(\beta x_i, \sigma_0^2 x_i^2).$$

Thus, the variance of  $Y_i$  is assumed to be proportional to  $x_i^2$ . In this exercise we will assume the value of  $\sigma_0^2$  to be known while the parameter  $\beta$  is estimated from the available data.

- a) Show that the estimator

$$\hat{\beta} = \frac{1}{n} \sum_{i=1}^n \frac{Y_i}{x_i}$$

is unbiased and find the variance of  $\hat{\beta}$ .

- b) What is the probability distribution of  $\hat{\beta}$ ? Give reason for the answer.

Work out a  $100(1 - \alpha)\%$ -confidence interval for  $\beta$ .

**Problem 4** The Botanical Research Station investigated how various kinds of fertilization affect the growth of sunflowers. Two kinds of fertilization were applied. Four randomly selected sunflowers were fertilized in one way, method A, and were  $x_i$  cm taller in a week, while four other randomly selected sunflowers were fertilized in another way, method B, and were  $y_i$  cm taller. Assume that  $x_i$  and  $y_i$  are independent observations from two normal distributions with the same variance. The growth gain of the plants were:

$x_i$ (method A)	12.0	10.2	12.1	10.4
$y_i$ (method B)	13.0	11.0	12.0	11.7

Det oppgis at  $\bar{x} = 11.175$ ,  $\sum(x_i - \bar{x})^2 = 3.0875$ ,  $\bar{y} = 11.925$ ,  $\sum(y_i - \bar{y})^2 = 2.0675$

- a) Perform a test to investigate whether method B gives a larger expected growth gains than method A. The null hypothesis is that A gives a growth gain at least as large as B gives. Use significance level  $\alpha = 0.05$ .
- b) Is  $p$ -value greater or less than 0.05?
- c) Find a 95% confidence interval for the difference between expected growth gains of the two fertilization methods.