



Løsningsforslag til ST1101 / ST6200 Sannsynlighetsregning  
Torsdag 7. juni 2007

**Oppgave 1**

$$\text{a) } E(Y) = \int_0^{\infty} y k e^{-y/\lambda} dy = k \int_0^{\infty} y^1 e^{-(1/\lambda)y} dy = \underline{\underline{k\lambda^2}}$$

$$E(Y^2) = k \int_0^{\infty} y^2 e^{-(1/\lambda)y} dy = \underline{\underline{2k\lambda^3}}$$

$$\text{Var}(Y) = E(Y^2) - (E(Y))^2 = 2k\lambda^3 - k^2\lambda^4 = \underline{\underline{k\lambda^3(2 - k\lambda)}}.$$

$$\text{b) } F_Y(y) = \int_0^y k e^{-y/\lambda} dy = \underline{\underline{k\lambda(1 - e^{-y/\lambda})}}, \text{ for } y \geq 0$$

$$P\left(\frac{1}{4}\mu \leq Y \leq \frac{3}{4}\mu\right) = F\left(\frac{3}{4}\mu\right) - F\left(\frac{1}{4}\mu\right) = k\lambda\left(1 - e^{-\frac{3\mu}{4\lambda}}\right) - k\lambda\left(1 - e^{-\frac{\mu}{4\lambda}}\right) = \underline{\underline{k\lambda\left(e^{-\frac{\mu}{4\lambda}} - e^{-\frac{3\mu}{4\lambda}}\right)}}$$

$$\text{c) } E(\bar{Y}) = E(Y) = \underline{\underline{k\lambda^2}}, \text{Var}(\bar{Y}) = \frac{1}{20} \text{Var}(Y) = \underline{\underline{\frac{k\lambda^3}{20}(2 - k\lambda)}}$$

$$\int_0^{\infty} k e^{-y/\lambda} dy = k\lambda = 1 \Rightarrow k = \underline{\underline{\frac{1}{\lambda}}}$$

$$P(\bar{Y} \geq 1.2\mu) = P\left(\frac{\bar{Y} - \lambda}{\lambda/\sqrt{20}} \geq \frac{1.2\lambda - \lambda}{\lambda/\sqrt{20}}\right) \approx P(z \geq 0.2\sqrt{20}) = P(z \geq 0.8944) \approx \underline{\underline{0.19}}$$

**Oppgave 2**

$$\text{a) } X \sim Po(\alpha/3) \Rightarrow E(X) = \alpha/3 = \underline{\underline{8/3}}$$

$$P(3 < X < 7) = P(X=4) + P(X=5) + P(X=6) = \frac{e^{-8/3} (8/3)^4}{4!} + \dots = \underline{\underline{0.26}}$$

$$\text{b) } Y \text{ er eksponensialfordelt med parameter } \lambda = 1/3.$$

$$T = 12 \times 5 + 12 \times Y \Rightarrow E(T) = 60 + 12 \times E(Y) = 60 + 12 \times 3 = \underline{\underline{96}}$$

$$\text{Var}(T) = 12^2 \text{Var}(Y) = 12^2 \times 3^2 \Rightarrow SD(T) = 12 \times 3 = \underline{\underline{36}}$$

c)  $H | x \sim \text{Bin}(x, p)$ ,  $E(H | x) = xp$

$$E(H) = E_x[E(H | x)] = E_x(xp) = pE_x(x) = \underline{\underline{\frac{p\alpha}{3}}}$$

$$f_H(h) = P(H = h) = \sum_x P(H = h, X = x) = \sum_x P(H = h | X = x)P(X = x)$$

$$= \sum_x \binom{x}{h} p^h (1-p)^{x-h} \frac{e^{-\alpha/3} (\alpha/3)^x}{x!} = \frac{e^{-\alpha/3} p^h (\alpha/3)^h}{h!} \sum_{x=h}^{\infty} \frac{[\alpha/3(1-p)]^{x-h}}{(x-h)!}$$

Bruker så  $y = x - h$  og at  $\sum_{y=0}^{\infty} \frac{a^y}{y!} = e^a$

$$\Rightarrow f_H(h) = \frac{e^{-p\alpha/3} (p\alpha/3)^h}{h!}. \text{ Dvs. Poissonfordeling med parameter } p\alpha/3.$$

### Oppgave 3

La  $A = \text{Dobbel } 6 \Rightarrow P(A) = \left(\frac{1}{6}\right)^2 = \frac{1}{36}$  og  $P(\bar{A}) = \frac{35}{36}$

Uavhengige kast  $\Rightarrow P(\text{Ikke } A \text{ i løpet av } n \text{ kast}) = \left(\frac{35}{36}\right)^n$

$$\Rightarrow P(\text{minst en } A \text{ i løpet av } n \text{ kast}) = 1 - \left(\frac{35}{36}\right)^n$$

Setter inn for  $n$  og finner at vi for  $n=24$  får 0.49, men for  $n=25$  får 0.506, så  $n=25$

### Oppgave 4

a)  $X \sim N(m, 0.01^2)$

$$m = 1.00 \Rightarrow P(X < 0.99) = P\left(\frac{X-1}{0.01} < \frac{0.99-1}{0.01}\right) = P(Z < -1) = \underline{\underline{0.1587}}$$

b)  $P(X < 0.99) = p = P(\text{undervektig})$

$$P(\text{ingen undervektige}) = (1-p)^{20}$$

$$P(\text{minst en undervektig}) = 1 - (1-p)^{20} < 0.01$$

$$(1-p)^{20} > 0.99 \Rightarrow p = 0.0005$$

Fra Tab. A1:  $\frac{0.99-m}{0.01} = -3.3 \Rightarrow m = \underline{\underline{1.023}}$