

PROBLEM SET 4

Let  $\mathcal{M}(\mathbb{R})$  be the family of moderate decrease functions. This means that  $f \in \mathcal{M}(\mathbb{R})$  if and only if  $f : \mathbb{R} \rightarrow \mathbb{C}$  is a continuous function, and there are constants  $\delta > 0$ ,  $M > 0$  and  $N > 0$  such that

$$|f(x)| \leq \frac{M}{|x|^{1+\delta}}$$

for  $|x| \geq N$ .

- (59) Let  $f : \mathbb{R} \rightarrow \mathbb{C}$  be a continuous function. Then,  $f \in \mathcal{M}(\mathbb{R})$  if and only if there is a constant  $K > 0$  such that

$$|f(x)| \leq \frac{K}{1 + |x|^{1+\delta}}$$

for all  $x \in \mathbb{R}$ .

- (60) Let  $\delta > 0$  be a real number. Then, for  $f \in \mathcal{M}(\mathbb{R})$ , we have

$$\delta \int_{-\infty}^{\infty} f(\delta x) dx = \int_{-\infty}^{\infty} f(x) dx.$$

- (61) Let  $f \in \mathcal{M}(\mathbb{R})$ .

- (a) Prove that  $\widehat{f}$  is an uniformly continuous function on  $\mathbb{R}$ .
- (b) Prove that  $\widehat{f}(\xi) \rightarrow 0$ , when  $\xi \rightarrow \pm\infty$  (Riemann-Lebesgue Lemma).
- (c) Show that if  $\widehat{f}(\xi) = 0$  for all  $\xi \in \mathbb{R}$ , then  $f$  is identically 0.

- (62) Suppose that  $F(x, y)$  is a continuous function in the plane  $(x, y) \in \mathbb{R}^2$ . Assume there is  $A > 0$  such that

$$|F(x, y)| \leq \frac{A}{(1 + |x|^{1+\delta})(1 + |y|^{1+\delta})}.$$

Define the functions

$$F_1(x) = \int_{-\infty}^{\infty} F(x, y) dy, \quad \text{and} \quad F_2(y) = \int_{-\infty}^{\infty} F(x, y) dx.$$

Therefore  $F_1, F_2 \in \mathcal{M}(\mathbb{R})$  and

$$\int_{-\infty}^{\infty} F_1(x, y) dx = \int_{-\infty}^{\infty} F_2(x, y) dy.$$

- (63) Compute the Fourier transform of the function

$$g(x) = \begin{cases} 1 + \cos x, & \text{if } |x| \leq \pi \\ 0, & \text{if } |x| > \pi. \end{cases}$$

(64) Define the function

$$f(x) = \begin{cases} 1 - |x|, & \text{if } |x| \leq 1 \\ 0, & \text{if } |x| > 1. \end{cases}$$

(a) Compute the Fourier transform of  $f$ .

(b) Determine the integral

$$\int_{-\infty}^{\infty} \left( \frac{\sin \pi \xi}{\pi \xi} \right)^4 d\xi.$$

(65) Let  $y > 0$ . Define the Poisson kernel as:

$$P_y(x) = \frac{1}{\pi} \frac{y}{x^2 + y^2},$$

Prove the following identities:

(a) For all  $x \in \mathbb{R}$ :

$$\int_{-\infty}^{\infty} e^{-2\pi|\xi|y} e^{2\pi i \xi x} d\xi = P_y(x).$$

(b) For all  $x \in \mathbb{R}$ :

$$\int_{-\infty}^{\infty} P_y(x) e^{-2\pi i \xi x} d\xi = e^{-2\pi|\xi|y}.$$

(66) Prove that the set  $\{P_y\}_{y>0}$  is a family of good kernels as  $y \rightarrow 0$ .

(67) Let  $a > 0$ . Compute the Fourier transform of the function

$$h_a(x) = \frac{a^2 - x^2}{(a^2 + x^2)^2}.$$

(68) Prove that if  $f \in \mathcal{M}(\mathbb{R})$  and

$$\int_{-\infty}^{\infty} f(y) e^{-y^2} e^{2xy} dy = 0,$$

for all  $x \in \mathbb{R}$ , then  $f = 0$ .

(69) The following exercise illustrates the principle that the decay of  $\widehat{f}$  is related to the continuity properties of  $f$ .

(a) Suppose that  $f \in \mathcal{M}(\mathbb{R})$  such that  $\widehat{f} \in \mathcal{M}(\mathbb{R})$ . Prove that  $f$  satisfies a Hölder condition of order  $\alpha$ , that is, that

$$|f(x+h) - f(x)| \leq M|h|^\alpha,$$

for some  $0 < \alpha < 1$ ,  $M > 0$  and  $x, h \in \mathbb{R}$ .

(b) Let  $f$  be a continuous function on  $\mathbb{R}$  which vanishes for  $|x| \geq 1$ , with  $f(0) = 0$ , and for  $0 < |x| < \delta$ ,

$$f(x) = \frac{1}{\ln(1/|x|)},$$

for some  $\delta > 0$ . Prove that  $\widehat{f} \notin \mathcal{M}(\mathbb{R})$ .

(70) Bump functions. Examples of compactly supported functions in  $\mathcal{S}(\mathbb{R})$  are very handy in many applications in analysis. Some examples are:

- (a) Suppose  $a < b$ , and  $f$  is the function such that  $f(x) = 0$  if  $x \leq a$  or  $x \geq b$  and

$$f(x) = e^{-1/(x-a)}e^{-1/(b-x)},$$

if  $a < x < b$ . Show that  $f$  is indefinitely differentiable on  $\mathbb{R}$ .

- (b) Prove that there exists an indefinitely differentiable function  $F$  on  $\mathbb{R}$  such that  $F(x) = 0$  if  $x \leq a$ ,  $F(x) = 1$  if  $x \geq b$ , and  $F$  is strictly increasing on  $[a, b]$ .
- (c) Let  $\delta > 0$  be so small such that  $a + \delta < b - \delta$ . Show that there exists an indefinitely differentiable function  $g$  such that  $g$  is 0 if  $x \leq a$  or  $x \geq b$ ,  $g$  is 1 on  $[a + \delta, b - \delta]$ , and  $g$  is strictly monotonic on  $[a, a + \delta]$  and  $[b - \delta, b]$ .

- (71) Let  $p, q \geq 1$  be real numbers,  $\alpha, \beta, m, n \in \mathbb{Z}$  be positive numbers. Suppose that there is a constant  $C > 0$  such that for any function  $G \in C^\infty(\mathbb{R})$  with compact support, the following inequality holds

$$\left( \int_{-\infty}^{\infty} \left| |\xi|^\alpha \frac{d^m \widehat{G}}{d\xi}(\xi) \right|^q d\xi \right)^{1/q} \leq C \left( \int_{-\infty}^{\infty} \left| |x|^\beta \frac{d^n G}{dx}(x) \right|^p dx \right)^{1/p}.$$

Give the necessary relationship between  $p, q, \alpha, \beta, m$  and  $n$ .

(Hint: Take a fixed function and its dilatations with  $\delta > 0$ . Taking  $\delta \rightarrow 0$  and  $\delta \rightarrow \infty$  one can obtain the desired relationship.)

*“How much better it is to get wisdom than gold,  
and to get knowledge is more to be desired than silver.”*

Email address: `carlos.a.c.chavez@ntnu.no`