



- 1 (a) We look at an interval $[x, x + \Delta x]$ on \mathbb{R} , and get that the conservation of mass in the interval is

$$\frac{\text{change of mass in the interval}}{\text{time}} = \text{flux in at } x - \text{flux out at } (x + \Delta x),$$
$$\Downarrow$$
$$\frac{d}{dt} \int_x^{x+\Delta x} \phi \rho(y, t) dy = j(x, t) - j(x + \Delta x, t).$$

Applying Leibniz's rule to the left hand side, we get

$$\int_x^{x+\Delta x} \phi \rho_t(y, t) dy = - \int_x^{x+\Delta x} j_x(y, t) dy.$$

In general, we have for a continuous function f that

$$\lim_{\Delta x \rightarrow 0} \left(\frac{1}{\Delta x} \int_{x_0}^{x_0+\Delta x} f(x) dx \right) = f(x_0),$$

and hence

$$\begin{aligned} \phi \rho_t(x, t) &= \lim_{\Delta x \rightarrow 0} \left(\frac{1}{\Delta x} \int_x^{x+\Delta x} \phi \rho_t(y, t) dy \right) \\ &= - \lim_{\Delta x \rightarrow 0} \left(\frac{1}{\Delta x} \int_x^{x+\Delta x} j_x(y, t) dy \right) = -j_x(x, t). \end{aligned}$$

Furthermore,

$$\begin{aligned} j_x(x, t) &= \frac{\partial}{\partial x} \left(-\rho \frac{k}{\mu} p_x \right) = \frac{\partial}{\partial x} \left(-\rho \frac{k}{\mu} \frac{\partial}{\partial x} (\rho RT) \right) \\ &= -\frac{kRT}{\mu} (\rho \rho_x)_x, \end{aligned}$$

and thus we have

$$\rho_t = K (\rho \rho_x)_x$$

with

$$K = \frac{kRT}{\mu \phi}.$$

(b) Since ρ_F is even and $\rho_F = 0$ for $x^2 > 12Ct^{\frac{2}{3}}$, we get

$$\begin{aligned} \int_{-\infty}^{\infty} \rho_F(x, t) dx &= 2 \int_0^{\sqrt{12Ct^{\frac{2}{3}}}} \rho_F(x, t) dx \\ &= 2 \int_0^{\sqrt{12Ct^{\frac{2}{3}}}} \left(Ct^{-\frac{1}{3}} - \frac{1}{12}x^2t^{-1} \right) dx \\ &= 2 \left[Ct^{-\frac{1}{3}}x - \frac{1}{12} \frac{1}{3}t^{-1}x^3 \right]_0^{\sqrt{12Ct^{\frac{2}{3}}}} \\ &= 2 \left(Ct^{-\frac{1}{3}}\sqrt{12Ct^{\frac{2}{3}}} - \frac{1}{12} \frac{1}{3}t^{-1}(12Ct^{\frac{2}{3}})^{\frac{3}{2}} \right) \\ &= 2 \left(\sqrt{12}C^{\frac{3}{2}} - \frac{1}{3}\sqrt{12}C^{\frac{3}{2}} \right) = \frac{4}{3}\sqrt{12}C^{\frac{3}{2}}. \end{aligned}$$

With $C = \frac{3^{\frac{1}{3}}}{4}$, we get

$$\int_{-\infty}^{\infty} \rho_F(x, t) dx = \frac{4}{3}\sqrt{12} \left(\frac{3^{\frac{1}{3}}}{4} \right)^{\frac{3}{2}} = \frac{8}{\sqrt{3}} \frac{\sqrt{3}}{8} = 1.$$

Following the hint and considering the regions separately, we immediately see that ρ_F satisfies the given equation in the regions $|x|^2 > 12Ct^{\frac{2}{3}}$. In the region $|x|^2 < 12Ct^{\frac{2}{3}}$, we have

$$\begin{aligned} (\rho_F)_t &= -\frac{1}{3}Ct^{-\frac{4}{3}} + \frac{1}{12}x^2t^{-2}, \\ (\rho_F)_x &= -\frac{1}{6}xt^{-1}, \\ (\rho_F)_{xx} &= -\frac{1}{6}t^{-1}, \end{aligned}$$

and hence

$$\begin{aligned} 2(\rho_F(\rho_F)_x)_x &= 2(\rho_F)_x^2 + 2\rho_F(\rho_F)_{xx} \\ &= 2\frac{1}{6^2}x^2t^{-2} + 2\left(-\frac{1}{6}Ct^{-\frac{1}{3}-1} + \frac{1}{6}\frac{1}{12}x^2t^{-2}\right) \\ &= \frac{1}{12}x^2t^{-2} - \frac{1}{3}Ct^{-\frac{4}{3}} \\ &= (\rho_F)_t, \end{aligned}$$

and thus the equation is satisfied by ρ_F also in this region.

(c) Because $\int_{-\infty}^{\infty} \rho_F dx = 1$, we have that

$$f(0) = \int_{-\infty}^{\infty} \rho_F(x, t)f(0) dx$$

and applying this and that $\rho_F \geq 0$, we get

$$\begin{aligned}
 \left| \int_{-\infty}^{\infty} \rho_F(x, t) f(x) dx - f(0) \right| &= \left| \int_{-\infty}^{\infty} \rho_F(x, t) (f(x) - f(0)) dx \right| \\
 &\leq \int_{-\infty}^{\infty} \rho_F(x, t) |f(x) - f(0)| dx \\
 &= \int_{-\sqrt{12Ct^{2/3}}}^{\sqrt{12Ct^{2/3}}} \rho_F(x, t) |f(x) - f(0)| dx \\
 &\leq \max_{|x| \leq \sqrt{12Ct^{2/3}}} |f(x) - f(0)| \cdot \int_{-\infty}^{\infty} \rho_F(x, t) dx \\
 &= \max_{|x| \leq \sqrt{12Ct^{2/3}}} |f(x) - f(0)|.
 \end{aligned}$$

Now we have

$$\lim_{t \rightarrow 0} \left| \int_{-\infty}^{\infty} \rho_F(x, t) f(x) dx - f(0) \right| \leq \lim_{\sqrt{12Ct^{2/3}} \rightarrow 0} \left(\max_{|x| \leq \sqrt{12Ct^{2/3}}} |f(x) - f(0)| \right) = 0,$$

since f is continuous, and $t \rightarrow 0$ implies $\sqrt{12Ct^{2/3}} \rightarrow 0$.

A fundamental solution is a solution with initial data $\rho_{F,0} = \delta_0$, where the delta function $\delta_0(x)$ is a function such that

$$\int_{-\infty}^{\infty} f(x) \delta_0(x) dx = f(0)$$

for any continuous function f . We have that

$$\int_{-\infty}^{\infty} \rho_F(x, 0) f(x) dx = \lim_{t \rightarrow 0} \int_{-\infty}^{\infty} \rho_F(x, t) f(x) dx = f(0),$$

and hence the initial solution $\rho_F(x, 0)$ of ρ_F is by definition a delta function.

- (d) The equation is given, and as initial solution we follow the hint and use a positive point source with integral 2. Thus we get

$$(1) \quad \begin{cases} h_t &= (h^2)_{xx}, \\ h(x, 0) &= 2\delta_0. \end{cases}$$

Following a similar deduction as when solving problem (b), we get that

$$\int_{-\infty}^{\infty} \rho_F(x, t) dx = 2 \quad \text{when } C = \left(\frac{3}{16} \right)^{\frac{1}{3}}.$$

We therefore have that equation (4) in the problem set is a solution to (1) above, with $C = \left(\frac{3}{16} \right)^{\frac{1}{3}}$. We then have that $h > 0$ for $|x|^2 < 12Ct^{\frac{2}{3}}$, and hence the extension of wet ground at $t = 10$ is given by

$$|x| = \sqrt{12Ct^{\frac{2}{3}}} = \sqrt{12 \left(\frac{3}{16} \right)^{\frac{1}{3}} 10^{\frac{2}{3}}} = 12^{\frac{1}{2}} \left(\frac{3}{16} \right)^{\frac{1}{6}} 10^{\frac{1}{3}} \approx 5.65.$$

- 2 (a) Denote by c the concentration of contaminants. There are two sources of flux. The first is the diffusive flux, given by Fick's law:

$$j_d = -\kappa \frac{\partial c}{\partial x}.$$

The second is the advective flux:

$$j_a = Uc.$$

Together, they form the total flux:

$$j = j_d + j_a = -\kappa \frac{\partial c}{\partial x} + Uc.$$

A point discharge will be carried a length L down the river after a time $T = L/U$. The discharge will spread out due to diffusion; the extent of this can be measured as in exercise 2c), i.e. the spread is proportional to $\sqrt{\kappa T} = \sqrt{\kappa L/U}$.

- (b) Using the flux from a) and noting that the conversion of substance A into B constitutes production terms, we state the conservation laws in integral form for an interval $[x_1, x_2]$ (we omit the t -dependence for readability):

$$\begin{aligned} \frac{d}{dt} \int_{x_1}^{x_2} a(x) dx &= \kappa \left(\frac{\partial a}{\partial x}(x_2) - \frac{\partial a}{\partial x}(x_1) \right) - U(a(x_2) - a(x_1)) - \int_{x_1}^{x_2} \mu a(x) dx, \\ \frac{d}{dt} \int_{x_1}^{x_2} b(x) dx &= \kappa \left(\frac{\partial b}{\partial x}(x_2) - \frac{\partial b}{\partial x}(x_1) \right) - U(b(x_2) - b(x_1)) + \int_{x_1}^{x_2} \mu a(x) - \lambda b(x) dx. \end{aligned}$$

To obtain the differential form of the conservation equation, we take $x_2 = x_1 + \Delta x$, divide by Δx and let $\Delta x \rightarrow 0$. This yields

$$\begin{aligned} \frac{\partial a}{\partial t} &= \kappa \frac{\partial^2 a}{\partial x^2} - U \frac{\partial a}{\partial x} - \mu a, \\ \frac{\partial b}{\partial t} &= \kappa \frac{\partial^2 b}{\partial x^2} - U \frac{\partial b}{\partial x} + \mu a - \lambda b. \end{aligned}$$

- (c) We now neglect diffusion and consider the convection-reaction equations:

$$\begin{aligned} \frac{\partial a}{\partial t} + U \frac{\partial a}{\partial x} &= -\mu a, & x > 0, t > 0, \\ \frac{\partial b}{\partial t} + U \frac{\partial b}{\partial x} &= \mu a - \lambda b & x > 0, t > 0. \end{aligned}$$

We need some boundary conditions. Since there is a constant rate of discharge of substance A at $x = 0$, we have a constant flux $j_A = Ua = q_0$ at $x = 0, t > 0$. Additionally, we may assume that the river is uncontaminated at $t = 0$, and that there is no discharge of substance B at $x = 0$. This gives the boundary conditions:

$$\begin{aligned} a(0, t) &= \frac{q_0}{U}, & t > 0 \\ a(x, 0) &= 0, & x > 0 \\ b(0, t) &= 0, & t > 0 \\ b(x, 0) &= 0, & x > 0. \end{aligned}$$

We solve the PDEs using the method of lines. Taking $z(t) = a(x(t), t)$, we get

$$\begin{aligned}\dot{z} &= -\mu z \\ \dot{x} &= U.\end{aligned}$$

Solving these ODEs yields

$$\begin{aligned}z(t) &= z(t_0)e^{\mu(t-t_0)} \\ x(t) &= U(t-t_0) + x_0.\end{aligned}$$

If $t_0 = 0$, we have $z(t_0) = 0$. Otherwise, $z(t_0) = \frac{q_0}{U}$. Now, to solve for $b(x, t)$, we take $w(t) = b(x(t), t)$, observing that the characteristics for a and b are identical. This gives us the ODE for w :

$$\begin{aligned}\dot{w} + \mu w &= \mu z \\ \Rightarrow \dot{w} + \mu w &= \frac{\mu q_0}{U} e^{\mu(t-t_0)},\end{aligned}$$

where we have disregarded the trivial case of characteristics starting at $t = 0$, which result in $b \equiv 0$. Now, using the hint, we obtain

$$w(t) = C_1 e^{\mu t} + \frac{\mu q_0}{U} t e^{\mu(t-t_0)},$$

and applying the initial condition $w(t_0) = 0$, we get

$$w(t) = (t - t_0) \frac{\mu q_0}{U} e^{\mu(t-t_0)},$$

yielding

$$b(U(t-t_0), t) = (t-t_0) \frac{\mu q_0}{U} e^{\mu(t-t_0)}.$$

We now "invert" by reintroducing $x = U(t-t_0)$ and get

$$b(x, t) = \begin{cases} x \frac{\mu q_0}{U^2} e^{\frac{\mu x}{U}}, & x < Ut \\ 0, & x > Ut. \end{cases}$$

To obtain the point of highest concentration of B , we fix a t and observe that, disregarding the case with $b = 0$:

$$\begin{aligned}\frac{d}{dx} b(x, t) &= \frac{\mu q_0}{U^2} e^{\frac{\mu x}{U}} \left(1 - \frac{\mu}{U} x\right) = 0 \\ \Rightarrow x &= \frac{U}{\mu}.\end{aligned}$$

This x is attainable if $t > \frac{1}{\mu}$. Otherwise, since $\frac{d}{dx} b(x, t) > 0$ for $x < \frac{U}{\mu}$, the maximum is attained at $x = Ut$.

- 3 (a) We do the computations with a segment with width B and introduce density, flux and sources. Here we assume that the density of sand ρ is a constant. The flux then becomes ρj and the source function will be $\rho q(x; t)$. (However, both B and ρ drop out from the relations at the end, such that we could as well compute per unit width, and with $\rho = 1$).

Our control volume has width B and extends from $x = x_0$ to $x = x_1$. Thus, we obtain the general (one-dimensional) conservation law

$$\frac{d}{dt} \int_{x_0}^{x_1} \rho B (b(x, t) - h) dx + \left(-k \frac{\partial b}{\partial x} (x_1, t) + k \frac{\partial b}{\partial x} (x_0, t) \right) \rho B = \int_{x_0}^{x_1} q(x, t) (\rho B) dx.$$

or

$$\frac{d}{dt} \int_{x_0}^{x_1} (b(x, t) - h) dx + \left(-k \frac{\partial b}{\partial x} (x_1, t) + k \frac{\partial b}{\partial x} (x_0, t) \right) = \int_{x_0}^{x_1} q(x, t) dx,$$

If we let $x_1 \rightarrow x_0$ and divide by $(x_1 - x_0)$, we obtain

$$\frac{\partial}{\partial t} (b - h) = \frac{\partial b}{\partial t} = k \frac{\partial^2 b}{\partial x^2} + q.$$

- (b) In this case, the source is localized at $x = 0$, such that the equation for $x > 0$ becomes just $b_t = kb_{xx}$. We scale b with h and the solution is

$$b = hf(x, t, k)$$

It is not obvious that we have a similarity solution since the depth h could be a length scale, but this length is not associated with the horizontal length. The problem is completely equivalent to a heat conduction problem where the temperature is constant and equal to T_0 at $x = 0$, and T_∞ when $x = \infty$. The temperature could then be written as $T(x, t) = T_0 + (T_\infty - T_0) \tau(x, t, k)$, and we obtain a similarity solution. Similarly to the temperature, we should be able to write the solution for b as

$$b = -h\beta \left(\frac{x}{\sqrt{kt}} \right) = -h\beta(\eta), \quad \eta = \frac{x}{\sqrt{kt}},$$

where $\beta(0) = 0$ and $\beta(\eta) \rightarrow 1$ when $\eta \rightarrow \infty$. Entering this into the equation after dividing by $-h$ lead to

$$\beta_t - k\beta_{xx} = -\frac{1}{2} \frac{x}{\sqrt{k}} \frac{1}{t^{3/2}} \beta' - k \frac{1}{kt} \beta'' = 0,$$

or

$$\beta'' + \frac{\eta}{2} \beta' = 0.$$

This is the equation given in the problem. By the hint and conditions $b(0, t) = 0$, $b(\infty, t) = -h$ we see that

$$b(x, t) = -h \operatorname{erf} \left(\frac{x}{\sqrt{kt}} \right).$$

- (c) The sand and clay volume at sea ($x \geq x_0$ here) is given by

$$\begin{aligned} V(t) &= \int_{x_0}^{\infty} B(h - b(x, t)) dx \\ &= \int_{x_0}^{Ut-x_0} B(h - b(x, t)) dx + \int_{Ut-x_0}^{\infty} B(h - b(x, t)) dx \\ &= BhUt + \int_0^{\infty} b_0(y) dy, \end{aligned}$$

where B was determined in a). By definition, $q_0 = V'(t)$, and hence

$$U = \frac{q_0}{Bh}.$$

If we let $\eta = x - Ut - x_0$ and put b into the equation where $x > Ut + x_0$, we obtain

$$-Ub'_0 = kb''_0.$$

thus

$$b''_0 + \frac{U}{k}b'_0 = 0,$$

with general solution

$$b_0(\eta) = C_1 + C_2 \exp\left(-\frac{U}{k}\eta\right).$$

It is required that

$$\begin{aligned} b_0(0) &= 0, \\ b_0(\infty) &= -h, \end{aligned}$$

such that the solution becomes

$$b_0(\eta) = h \left(\exp\left(-\frac{U}{k}\eta\right) - 1 \right).$$

Introducing the original variables leads to

$$b(x, t) = \begin{cases} 0, & x \leq s(t) = Ut + x_0, \\ h \left(\exp\left(-\frac{U}{k}(x - Ut - x_0)\right) - 1 \right), & x > Ut + x_0. \end{cases}$$