TMA4267 Linear Statistical Models V2017 (L17)

Part 4: Design of Experiments

Mette Langaas

Department of Mathematical Sciences, NTNU

To be lectured: March 21, 2017

1/16

Design of experiments vs. observational studies

In this part of the course we are working with the linear regression model:

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon} \text{ with } \boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma^2 \mathbf{I})$$

and use results from Part 2 of the course.

Earlier in the course: both the design matrix X and the reponses Y were observed together in a randomly selected sample from a population.

- Munich rent index: rent prices vs. area, location, condition of bathroom, condition of kitchen,
- Lakes: pH level vs. content of SO₄, NO₃, latent Al, Ca, organic, position, area.
- ▶ Happiness: Happiness vs. love, money, sex and work.

Now: we choose (design) the experiment by specifying the design matrix \boldsymbol{X} to be used to produce a sample, and then collecting reponses Y for this design matrix.

2/16

Today:

- ▶ Observational studies vs. designed experiments.
- ► Still linear regression, but now with *k* factors each with only 2 levels.
- ▶ Effect coding, orthogonal columns in design matrix.
- ▶ 2^k full factorial design.
- ▶ Simplified formulas for $\hat{\beta}$, $Cov(\hat{\beta})$ and SSE.
- ► If time: from parameter estimated to main and interaction effects.

Part 4 is based on Tyssedal: Design of experiments note.

1/16

The pilot plant example - Version 1

At a pilot plant a chemical process is investigated.

- ► The outcome of the process is measured as chemical yield (in grams).
- ► Two quantitative variables (factors) were investigated:
 - ► Factor A: Temperature (in degrees C).
 - ► Factor B: Concentration (in percentage).

Experiment no.	Temperature	Concentration	Yield
1	160	20	60
2	180	20	72
3	160	40	54
4	180	40	68
	<i>x</i> ₁	<i>x</i> ₂	у

Regression with pilot plant data V1- original

```
> x1=c(160,180,160,180)
> x2=c(20,20,40,40)
> y=c(60,72,54,68)
> fitx=lm(v~x1*x2)
Coefficients:
(Intercept)
                     x1
                                  x2
                                            x1:x2
    -14.000
                  0.500
                              -1.100
                                            0.005
> model.matrix(fitx)
  (Intercept) x1 x2 x1:x2
           1 160 20 3200
           1 180 20 3600
           1 160 40 6400
           1 180 40 7200
```

4/16

Regression with original and coded factors

Original: x_1 and x_2 , gave estimated regression equation

$$\hat{\mathbf{y}} = -14 + 0.5\mathbf{x}_1 - 1.1\mathbf{x}_2 + 0.005\mathbf{x}_1 \cdot \mathbf{x}_2$$

Coded: $z_1 = (x_1 - 170)/10$ and $z_2 = (x_2 - 30)/10$, gave estimated regression equation

$$\hat{\mathbf{y}} = 63.5 + 6.5z_1 - 2.5z_2 + 0.5z_1 \cdot z_2$$

Can you compare these two results?

6/16

Regression with pilot plant data V1- recoded

```
> # recode to -1 and 1
> z1=(x1-(max(x1)+min(x1))/2)/((max(x1)-min(x1))/2)
> z2=(x2-(max(x2)+min(x2))/2)/((max(x2)-min(x2))/2)
> fitz=lm(v^z1*z2)
Coefficients:
(Intercept)
                     z1
                                           z1:z2
      63.5
                    6.5
                                -2.5
                                             0.5
> model.matrix(fitz)
  (Intercept) z1 z2 z1:z2
           1 -1 -1
           1 1 -1 -1
           1 -1 1 -1
           1 1 1
```

5/16

Regression with original and coded factors

Substitute $z_1 = (x_1 - 170)/10$ and $z_2 = (x_2 - 30)/10$ into the equation to get a estimated regression equation based on x_1 and x_2 .

$$\hat{y} = 63.5 + 6.5z_1 - 2.5z_2 + 0.5z_1 \cdot z_2$$

$$= 63.5 + 6.5\frac{x_1 - 170}{10} - 2.5\frac{x_2 - 30}{10} + 0.5\frac{x_1 - 170}{10} \cdot \frac{x_2 - 30}{10}$$

$$= 63.5 - 6.5\frac{170}{10} + 2.5\frac{30}{10} + 0.5\frac{170 \cdot 30}{10 \cdot 10}$$

$$+ x_1(6.5\frac{1}{10} - 0.5\frac{1}{10}\frac{30}{10}) + x_2(-2.5\frac{1}{10} - 0.5\frac{1}{10}\frac{170}{10})$$

$$+ 0.5\frac{1}{10}\frac{1}{10}x_1 \cdot x_2$$

$$= -14 + 0.5x_1 - 1.1x_2 + 0.005x_1 \cdot x_2$$

Design of experiments (DOE) terminology

- ▶ Variables are called factors, and denoted A, B, C, ...
- ▶ We will only look at factors with two levels:
 - ▶ high, coded as +1 or just +, and,
 - ▶ low, coded as −1 or just −.
- ▶ In the pilot plant example we had two factors with two levels, thus 2 · 2 = 4 possible combinations. In general *k* factors with two levels gives 2^k possible combinations.

Standard notation for 2² experiment:

Experiment no.	Α	В	AB	Level code	Response
1	-1	-1	1	1	<i>y</i> ₁
2	1	-1	-1	а	<i>y</i> ₂
3	-1	1	-1	Ь	<i>y</i> 3
4	1	1	1	ab	<i>y</i> 4
	z_1	z_2	<i>z</i> ₁₂		у

8/16

Main effects in DOE

Main effect of A

$$\hat{A} = 2\hat{\beta}_1$$

$$= \frac{y_2 + y_4 + y_6 + y_8}{4} - \frac{y_1 + y_3 + y_5 + y_7}{4}$$

Interpretation: mean response when \boldsymbol{A} is high MINUS mean response when \boldsymbol{A} is low.

Similarly, main effect of B

$$\hat{B} = 2\hat{\beta}_2$$

$$= \frac{y_3 + y_4 + y_7 + y_8}{4} - \frac{y_1 + y_2 + y_5 + y_6}{4}$$

Interpretation: mean response when ${\it B}$ is high MINUS mean response when ${\it B}$ is low.

10 / 16


Lima beans example

Experiment from Box, Hunter, Hunter, Statistics for Experimenters, page 321.

- ► A: depth of planting (0.5 inch or 1.5 inch)
- ▶ B: watering daily (once or twice)
- ► C: type of lima bean (baby or large)
- Y: yield

Α	В	С	AB	AC	BC	ABC	Level code	Response
-	-	-	+	+	+	-	1	6
+	-	-	-	-	+	+	a	4
-	+	-	-	+	-	+	ь	10
+	+	-	+	-	-	-	ab	7
-	-	+	+	-	-	+	c	4
+	-	+	-	+	-	-	ac	3
- 1	+	+	-	-	+	-	bc	8
+	+	+	+	+	+	+	abc	5
×1	X2	Х3	X12	X13	X23	X123		y

9/16



Explain the main effects in plain words!

A: depth (0.5 or 1), B: watering daily (once, twice), C: type (baby, large).

Interaction effect in DOE

- ▶ What is the terpretation in DOE associated with β_{12} ?
- ▶ In DOE $2\hat{\beta}_{12}$ is denoted \widehat{AB} and is called the *estimated* interaction effect between A and B.

$$\widehat{AB} = 2\widehat{\beta}_{12}$$

$$= \frac{\text{estimated main effect of } A \text{ when } B \text{ is high}}{2}$$

$$- \frac{\text{estimated main effect of } A \text{ when } B \text{ is low}}{2}$$

$$= \frac{\text{estimated main effect of } B \text{ when } A \text{ is high}}{2}$$


$$- \frac{\text{estimated main effect of } B \text{ when } A \text{ is low}}{2}$$

12 / 16


Interpretation of \widehat{ABC}

- ▶ $\widehat{ABC} = \frac{1}{2}\widehat{AB}$ interaction when C is at the high level $\frac{1}{2}\widehat{AB}$ interaction when C is at the low level.
- Or, two other possible interpretation with swapped placed for A, B and C.
- ▶ And remember that $\widehat{AB} = \frac{1}{2}\widehat{A}$ main effect when B is at the high level $\frac{1}{2}\widehat{A}$ main effect when B is at the low level.

14/16



Geometric interpretation of effects



2^k full factorial

- ▶ There are k factors: A, B, C, ..., and
- ▶ 2=each factor has two levels.
- ▶ There are 2^k possible experiments.
- \blacktriangleright We have in total 2^k parameters to be estimated:
 - ▶ 1 intercept
 - $k = \binom{k}{1}$ main effects: A, B, C, ...
 - (k) two factor interactions: AB, AC, ..., BC, BD,...
 - $\binom{\overline{k}}{3}$ three factor interactions: ABC, ABD, ABE, ...
 - **>** ...
 - $\binom{k}{k} = 1$ k factor interaction.

$$Y_{i} = \beta_{0} + \beta_{1}x_{1i} + \beta_{2}x_{2i} + \dots + \beta_{k}x_{ki}$$

$$+ \beta_{12}x_{12} + \dots + \beta_{k-1,k}x_{k-1,k}$$

$$+ \beta_{123}x_{123} + \dots + \beta_{k-2,k-1,k}x_{k-2,k-1,k}$$

$$\dots + \beta_{12\dots k}x_{12\dots k}$$

16 / 16

Observe that each foctor color has $\frac{\eta}{1+\epsilon}\chi_j=0$, and we also include an vibraget tom with $\frac{\eta}{1+\epsilon}\chi_{ij}=n$ \Longrightarrow χ_{ij}

& and 55K will howesuple formulae of this fell 2 dation

2

```
THAY267 LIA
Port 4: Design of Exporments
                                     21.08.2017
with 2 factorial designs
Regression Y= Xp+c , E~Nn(0, +T)
                  thep, intropt and it owners
 B= (XTX)-1 XTY ~ Np ( p, (XTX)-1) + nove!
 And we used observelingal data
Now: we design the experiment = Choose X!
 How should we choose X? Achieve some kind of
ophnality.
    - unvinise ( ( ) = fr ( 2, ( XIX)-1)
    - minimize det (Cor(67))
 Our focus:
    - maximize interpretability; e.g. by showing X
     so that Cov(f), fu)=0 ( (BTE) is diagonal
      which we may advisite by choosing the column of
     I to be orthogonal to ead other.
     We focus on 2k factorid design look at k factor each at 2 Course
```


3

Hands-on:

- 1) Show that say two columns of X ere orthogonal, " Xij Xik = 0
- 2) Show that $\sum_{i=1}^{n} X_{ij} = 0$ for all j except j=0
- 3) And that \(\frac{1}{2}\xi_j^2 = n.\)

Now: How does this (1+2+9) influence our formulas for i) if ii) SSR - \(\hat{\text{L}} \) (ii) SSR - \(\hat{\text{L}} \) (ij)

$$\hat{\beta} = \begin{bmatrix} \frac{1}{12} & 0 & 0 \\ 0 & \frac{1}{12} & 0 \end{bmatrix} \times \begin{bmatrix} \frac{1}{12} & \frac{1}{12} & \frac{1}{12} & \frac{1}{12} \\ \frac{1}{12} & \frac{1}{12} & \frac{1}{12} & \frac{1}{12} \end{bmatrix}$$

$$SSR = \sum_{i=1}^{n} ((\hat{y}_{i} - \hat{y}_{j})^{n} = \sum_{i=1}^{n} (\sum_{j=0}^{n-1} \hat{p}_{j} x_{ij} - \hat{p}_{i})^{2}$$

$$= \sum_{i=1}^{n} (\hat{p}_{0}^{h} + \sum_{j=1}^{n-1} \hat{p}_{j} x_{ij} - \hat{p}_{0}^{h})^{2} = \sum_{i=1}^{n} (\sum_{j=1}^{n-1} \hat{p}_{j} x_{ij})^{2}$$

$$= \sum_{i=1}^{n} (\hat{p}_{0}^{h} + x_{ii} + \hat{p}_{0}^{h} + x_{ii} + p_{0}^{h} + p_{0}^{$$

Observe that
$$\hat{F}_{ij}$$
 is only dependent on \hat{X}_{ij} and not on \hat{X}_{ik} has \hat{F}_{ij} is only dependent on \hat{X}_{ij} and not on \hat{X}_{ik} has \hat{F}_{ij} so \hat{F}_{ij} will not drow if we drow that nearly. If we drow the transfer the nearly \hat{F}_{ij} of $\hat{F$

TMA4267 Linear Statistical Models V2017 (L18)

Part 4: Design of Experiments Full 2^k factorial designs DOE Effects, estimating variability and performing inference Compulsory DOE project

Mette Langaas

Department of Mathematical Sciences, NTNU

To be lectured: March 24, 2017

Last lesson - and today:

- ▶ Observational studies vs. designed experiments.
- ▶ Still linear regression, but now with *k* factors each with only 2 levels.
- ▶ Effect coding, orthogonal columns in design matrix.
- ▶ 2^k full factorial design.
- ▶ Simplified formulas for $\hat{\beta}$, $Cov(\hat{\beta})$ and SSE.
- ► From parameter estimated to main and interaction effects.
- ► Inference.
- ► Compulsory exercise 4: the DOE project

Part 4 is based on Tyssedal: Design of experiments note.

1 / 45

Design of experiments (DOE) terminology

- ▶ Variables are called factors, and denoted A, B, C, ...
- ▶ We will only look at factors with two levels:
 - ▶ high, coded as +1 or just +, and,
 - ▶ low, coded as −1 or just −.
- ▶ The lima beans example had three factors with two levels, thus $2^3 = 8$ possible combinations. In general k factors with two levels gives 2^k possible combinations.

Standard notation for 2³ experiment (responses for lima beans included)

Α	В	С	AB	AC	BC	ABC	Level code	Response
-	-	-	+	+	+	-	1	6
+	-	- 1	-	-	+	+	a	4
-	+	- 1	-	+	-	+	ь	10
+	+	-	+	-	-	-	ab	7
-	-	+	+	-	-	+	c	4
+	-	+	-	+	-	-	ac	3
-	+	+	-	-	+	-	bc	8
+	+	+	+	+	+	+	abc	5
×1	× ₂	×3	×12	×13	X23	×123		y

3 / 45

Lima beans example

Experiment from Box, Hunter, Hunter, Statistics for Experimenters, page 321.

- ► A: depth of planting (0.5 inch or 1.5 inch)
- ► B: watering daily (once or twice)
- ► C: type of lima bean (baby or large)
- Y: yield

Research question: what is the combination of A, B, C giving the highest vield?

2/45

Results from last lecture: 2^k full factorial

Known from Part 2: $\hat{\boldsymbol{\beta}} = (\boldsymbol{X}^T \boldsymbol{X})^{-1} \boldsymbol{X}^T \boldsymbol{Y}$ and $Cov(\hat{\boldsymbol{\beta}}) = \sigma^2(\boldsymbol{X}^T\boldsymbol{X})^{-1}.$

- ▶ The design matrix is chosen so that the columns (containing -1 and 1) are orthogonal, and thus
 - $\sum_{i=1}^{n} x_{ij} x_{ik} = 0$ for all combinations of the columns of the design matrix X.
 - $\sum_{i=1}^{n} x_{ii}^2 = n.$
- ▶ The orthogonal columns lead to that the following formulas are easy to interpret and calculate:
 - $\boldsymbol{X}^T \boldsymbol{X} = \text{diagonal matrix with } n \text{ on the diagonal.}$
 - $\hat{\beta}_j = \frac{1}{n} \sum_{i=1}^n x_{ij} Y_i.$

 - $Var(\hat{\beta}_j) = \frac{\sigma^2}{n}.$ $Vov(\hat{\beta}_j, \hat{\beta}_k) = 0 \text{ for all } j \neq k.$
 - ► SSR= $\sum_{i=1}^{p-1} \hat{\beta}_i^2$.

See class notes for L17 for details on the derivation.


Lima beans example: full 2³ factorial design

- ► A: depth of planting (0.5 inch or 1.5 inch)
- ► B: watering daily (once or twice)
- ► C: type of lima bean (baby or large)
- Y: yield

Α	В	C	AB	AC	BC	ABC	Level code	Response
-	-	-	+	+	+	-	1	6
+	-	-	-	-	+	+	а	4
-	+	-	-	+	-	+	Ь	10
+	+	-	+	-	-	-	ab	7
-	-	+	+	-	-	+	С	4
+	-	+	-	+	-	-	ac	3
-	+	+	-	-	+	-	bc	8
+	+	+	+	+	+	+	abc	5
<i>x</i> ₁	<i>x</i> ₂	<i>X</i> 3	X ₁₂	X ₁₃	X23	X ₁₂₃		у

Write down the regression model with all possible interactions, and find $\hat{\beta}_i = \frac{1}{n} \sum_{i=1}^n x_{ij} Y_i$ for the A and the AB columns.

5 / 45



A B C A:B A:C B:C A:B:C -2.25 3.25 -1.75 -0.75 0.25 -0.25 -0.25

Explain the main effects in plain words!

A: depth (0.5 or 1), B: watering daily (once, twice), C: type (baby, large).

7 / 45

Main effects in DOE

Main effect of A

$$\hat{A} = 2\hat{\beta}_1$$

$$= \frac{y_2 + y_4 + y_6 + y_8}{4} - \frac{y_1 + y_3 + y_5 + y_7}{4}$$

Interpretation: mean response when A is high MINUS mean response when A is low.

Similarly, main effect of B

$$\hat{B} = 2\hat{\beta}_2
= \frac{y_3 + y_4 + y_7 + y_8}{4} - \frac{y_1 + y_2 + y_5 + y_6}{4}$$

Interpretation: mean response when B is high MINUS mean response when B is low.

6 / 45

Interaction effect in DOE

- ▶ What is the terpretation in DOE associated with β_{12} ?
- ▶ In DOE $2\hat{\beta}_{12}$ is denoted \widehat{AB} and is called the *estimated* interaction effect between A and B.


$$\widehat{AB} = 2\widehat{\beta}_{12}$$

$$= \frac{\text{estimated main effect of } A \text{ when } B \text{ is high}}{2}$$

$$- \frac{\text{estimated main effect of } A \text{ when } B \text{ is low}}{2}$$

$$= \frac{\text{estimated main effect of } B \text{ when } A \text{ is high}}{2}$$

$$- \frac{\text{estimated main effect of } B \text{ when } A \text{ is low}}{2}$$



9 / 45

R: DOE set-up for lima beans

```
> library(FrF2)
> plan <- FrF2(nruns=8,nfactors=3,randomize=FALSE)
creating full factorial with 8 runs ...
> plan
```

A B C

1 -1 -1 -1

2 1 -1 -:

3 -1 1 -1

4 1 1 -1

5 -1 -1 1

6 1 -1 1

7 -1 1 1 8 1 1 1

class=design, type= full factorial

But, the experiment should be performed in *random order*. We use R to find the random order, and then we choose randomize=TRUE. I have used randomize=FALSE here because the y-values were easier to read in in standard order.

11 / 45

Interpretation of \widehat{ABC}

- ▶ $\widehat{ABC} = \frac{1}{2}\widehat{AB}$ interaction when C is at the high level $\frac{1}{2}\widehat{AB}$ interaction when C is at the low level.
- Or, two other possible interpretation with swapped placed for A B and C
- ▶ And remember that $\widehat{AB} = \frac{1}{2}\widehat{A}$ main effect when B is at the high level $\frac{1}{2}\widehat{A}$ main effect when B is at the low level.

10 / 45

R: DOE add response

```
> y <- c(6,4,10,7,4,3,8,5)
> y
[1] 6 4 10 7 4 3 8 5
> plan <- add.response(plan,y)
> plan
    A B C y
1 -1 -1 -1 6
2 1 -1 -1 4
3 -1 1 -1 10
4 1 1 -1 7
5 -1 -1 1 4
6 1 -1 1 3
7 -1 1 1 8
8 1 1 1 5
class=design, type= full factorial
```


R: DOE Im and effect

- > lm3 <- lm(y~(.)^3,data=plan)
- > MEPlot(1m3)
- > IAPlot(1m3)
- > effects <- 2*lm3\$coeff
- > effects

(Intercept) A1 B1 C1 A1:B1 A1:C1 B1:C1 A1:B1:C1 11.75 -2.25 3.25 -1.75 -0.75 0.25 -0.25 -0.25

13 / 45

Geometric interpretation of effects



15 / 45

2^k full factorial

- ► There are k factors: A, B, C, ..., and
- ▶ 2=each factor has two levels.
- ▶ There are 2^k possible experiments.
- \blacktriangleright We have in total 2^k parameters to be estimated:
 - ▶ 1 intercept
 - $k = \binom{k}{1}$ main effects: A, B, C, ...
 - (a) two factor interactions: AB, AC, ..., BC, BD,...
 - $\binom{\overline{k}}{3}$ three factor interactions: ABC, ABD, ABE, ...
 - **•** • •
 - $\binom{k}{k} = 1$ k factor interaction.

$$Y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k$$

$$+ \beta_{12} x_{12} + \dots + \beta_{k-1,k} x_{k-1,k}$$

$$+ \beta_{123} x_{123} + \dots + \beta_{k-2,k-1,k} x_{k-2,k-1,k}$$

$$\dots + \beta_{12...k} x_{12...k} + \varepsilon$$

14 / 45

Lima beans: significant effects?

Lima beans: significant effects?

17 / 45


Three factors in three full replicates


- ▶ Lima beans experiment from Box, Hunter, Hunter page 321.
 - ► A: depth of planting (0.5 inch or 1.5 inch)
 - ▶ B: watering daily (once or twice)
 - ► C: type of limabean (baby or large)
 - Y: yield
- ► r = 3: Performed in three full replicate experiments, i.e. three measurements for each combination of A, B and C.
- ▶ We then have $(r-1)2^3 = 2 \cdot 8 = 16$ degrees of freedom for estimating the error variance.
- Estimates follow automatically. Perform this for yourself. R code on course www-page.

19 / 45

Estimation of σ^2

- 1. Perform replicates, estimate the full model and use s^2 from regression model.
- 2. Assuming specified higher order interactions are zero (changing the regression model).
- 3. If the two above is not possible: Lenth's Pseudo Standard Error (PSE).





Back to no extra replicates: Lima beans with only main effects

ANOVA output: R

Analysis of Variance Table

```
Response: y
         Df Sum Sq Mean Sq F value
          1 28.167 28.167 52.0000 2.075e-06 ***
          1 37.500 37.500 69.2308 3.319e-07 ***
C
          1 24.000 24.000 44.3077 5.517e-06 ***
A:B
          1 0.667
                     0.667 1.2308
          1 0.167
                     0.167 0.3077
                                     0.5868
          1 0.167
                     0.167 0.3077
                                     0.5868
A:B:C
                                     1.0000
          1 0.000
                     0.000
                           0.0000
```

0.542

22 / 45

Back to no extra replicates: Assuming specified higher order interactions are zero
Result that is JUST a curiosity

In general

Residuals 16 8.667

$$\widehat{\textit{Effect}}_j \sim \textit{N}(\textit{Effect}_j, \sigma_{\textit{effect}}^2)$$

▶ If we assume that the effect is zero ($\beta_j = 0$), then $E(\textit{Effect}_i) = 0$ and

$$E(\widehat{Effect}_{i}^{2}) = \sigma_{effect}^{2}$$

- ▶ Thus $\widehat{\it Effect}_i^2$ is an unbiased estimator of $\sigma_{\it effect}^2$ if $\beta_j = 0$.
- ▶ If several effects are assumed to be 0, we use the average of the $\widehat{\it Effect}_{j}^{2}$ to estimate $\sigma_{\it effect}^{2}$.

Lima beans estimated effects: full model

```
Estimated effects (2*coeff): (Intercept) At B1 C1 A1:B1 A1:C1 B1:C1 A1:B1:C1 I1.75 -2.25 3.25 -1.75 -0.75 0.25 -0.25 -0.25 -0.25 Analysis of Variance Table

Df Sum Sq Hean Sq F value Pr(>F)

A 1 10.125 10.125

B 1 21.125 21.125

C B 1 6.125 6.125

A:B 1 1.125 1.125

A:C 1 0.125 0.125


B:C 1 0.125 0.125

B:C 1 0.125 0.125

A:B:C 1 0.125 0.125
```

25 / 45

R: Pareto plot for Lima beans



Pareto plot: ordered histogram of absolute value of estimated effects, Length sign line added.

27 / 45

Lenth's PSE

Let C_1, C_2, \ldots, C_m be estimated effects, e.g. $\hat{A}, \hat{B}, \widehat{AB}$, etc.

- 1. Order absolute values $|C_i|$ in increasing order.
- 2. Find the median of the $|C_i|$ and compute preliminary estimate

$$s_0 = 1.5 \cdot \mathsf{median}_j |C_j|$$

3. Take out the effects C_j with $|C_j| \ge 2.5 \cdot s_0$ and find the median of the rest of the $|C_j|$. Then PSE is this median multiplied by 1.5, i.e.

$$PSE = 1.5 \cdot median\{|C_i| : |C_i| < 2.5s_0\}$$

and this is Lenth's estimate of $\sigma_{\it effect}$.

4. Lenth has suggested empirically that the degrees of freedom to be used with PSE is m/3 where m is the initial number of effects in the algorithm (intercept not included). Thus we claim as significant the effects for which $|C_j| > t_{\alpha/2, m/3} \cdot PSE$.

26 / 45

Which ν ?

From the previous slide, connection between ν and your chosen estimation method for σ and $\sigma_{\it effect}$.

- 1. If you have performed the 2^k experiment r times, then $\nu=(r-1)2^k$.
- 2. If m effects (preferrable higher order interactions) are assumed to be zero, then $\nu=m$.
- 3. When Lenth's PSE is used, the degrees of freedom is

$$\nu = \frac{2^k - 1}{3}$$

where $2^k - 1$ is the number of effects in the model, while the 3 in the denominator has been found empirically by Lenth.

DOE workflow

- 1. Set up full factorial design with k factors in R, and
- 2. randomize the runs.
- 3. Perform experiments, and enter data into R.
- 4. Fit a full model (all interactions) make Pareto-plot (with/without red line).
- If you do not have replications, refit the data to a reduced model.
- 6. Assess model fit (residual plots, need transformations?).
- 7. Construct confidence intervals, assess significance.
- 8. Interpret you results (main and interaction plots).

29 / 45



Response: length of plant after 8 days of growing.

31 / 45

Example compulsory project

"From a seed to a nice plant"



30 / 45

The experiments

StdOrder	RunOrder	CenterPt	Blocks	Seeds	Watering fluid		Additional nutrients	Length (response variable)
5	1	1	1	-1	-1	1	-1	0.1
2	2	1	1	1	-1	-1	-1	20.3
16	3	1	1	1	1	1	1	0.9
9	4	1	1	-1	-1	-1	1	0.2
15	5	1	1	-1	1	1	1	0.0
12	6	1	1	1	1	-1	1	6.9
6	7	1	1	1	-1	1	-1	1.1
1	- 8	1	1	-1	-1	-1	-1	11.7
10	9	1	1	1	-1	-1	1	5.9
13	10	1	1	-1	-1	- 1	1	0.0
4	11	1	. 1	1	1	-1	-1	23.3
8	12	1	1	1	1	1	-1	4.5
7	13	1	1	-1	1	1	-1	9.1
3	14	1	1	-1	1	-1	-1	12.2
14	15	1	1	1	-1	1	1	1.5
11	16	1	1	-1	1	-1	1	2.9

Full model


Estimated Effects and Coefficients for length (coded units)

Term	Effect	Coef
Constant		6,287
A	3,525	1,763
В	2,375	1,187
C	-8,275	-4,138
D	-8,000	-4,000
A*B	-0,675	-0,337
A*C	-3,825	-1,913
A*D	-0,500	-0,250
B*C	0,575	0,287
B*D	-1,600	-0,800
C*D	4,900	2,450
A*B*C	-0,875	-0,438
A*B*D	0,100	0,050
A*C*D	2,000	1,000
B*C*D	-1,650	-0,825
A*B*C*D	1,150	0,575

33 / 45

35 / 45

Inference










Figure 5.7 Normal plot of the effects with terms up to 2nd order.

A, C and D, AC and CD found to be significant.

Full model










Figure 5.3 Normal plot of the effects with terms up to 4th order.

34 / 45

Interpretation: Interaction plots








Figure 6.2 Interaction plot between seeds and growth medium (AC).

The practical issues (1)

- ► You may work alone, or in groups of two.
- ► You need to perform a multiple regression experiment consisting of 16 trials - that is, n=16 observations.
- ▶ The response that is measure should be continuous, so that the response itself or a transformation of the response in a regression model can be seen to be normally distributed. (It is also possible to assume that a response with at least 7 ordered categories can be seen as continuous.)
- ▶ You choose 3 or 4 factors with two levels each that might influence your response (it is possible to choose more factors, but then you need to do a so called fractional factorial design to be lectured soon).

37 / 45

Genuine run replicates

"When genuine run replicates are made under a given set of experimental conditions, the variation between the associated observations may be used to estimate the standard deviation of the effects. By genuine run replicated we mean that variation between runs made at the same experimental conditions is a reflection of the total variability afflicting runs made at different experimental conditions. This point requires careful consideration."

39 / 45

The practical issues (2)

- ▶ If you choose 3 factors you need to perform all possible combinations of the 3 factors two times (2.2.2=8), if you choose 4 factors you need to perform all possible combinations only once $(2 \cdot 2 \cdot 2 \cdot 2 = 16)$. If you choose more than 4 factors you need to study the "factional factorials" to find out which of the possible combinations you perform.
- ▶ A very important aspect of performing the 16 trials is that the trials should be independent and performed in a randomized order (why?). You use R to randomize the experiments for you.
- ► Each experiment should be a complete new experiment a genuine run replicate, unless you use blocking (not lectured yet). For example a block effect my be person or day.

38 / 45

Genuine run replicates

Randomization of run order usually ensures that replicates are genuine. Pilot plant example: each run consists of

- 1. cleaning the reactor
- 2. inserting the appropriate catalyst carge
- 3. running the apparatus at at given temperature and a given feed concentration for 3 hrs to allow the process to settle down at the chosen experimental conditions, and
- 4. combining chemical analyses made on these samples.

A genuine run replicate must involve the taking of all these steps again. In particular, several chemical analyses from a single run would provide only an estimate of analytical variance, usually only a small part of the run-to-run variance. From Box, Hunter, Hunter (1978, 2005): "Statistics for Experimenters", Ch.10.6

The practical issues (3)

- After you have performed all 16 experiments you need to record the response and enter it into the experiment you have designed in R.
- ► Then you analyze the data, estimate effects, perform inference, check the model assumptions (RESIDUALS!), and explain your findings.

41 / 45

The report (2)

- 4. Choice of design: 2 k factorial, 2 k-p fractional factorial (resolution?)? Is it necessary or desirable to use a blocked design? Is it necessary or desirable with replicates?
- 5. Implementation of the experiment: Randomization. Describe any problems with the implementation.
- 6. Analysis of data: Calculation of effects and assessment of statistical significance. Use Lenth (not only), replicates or "setting some interactions to zero" to perform inference? Check the assumptions. RESIDUAL PLOTS!
- 7. Conclusion (explain main and interaction plots) and recommendations: Which conclusions can you draw from the experiment?

To get 10 points you need to have addressed all of these aspects in a correct manner! BUT - don't hand in more than 8 pages (included printout from R and plots)!

43 / 45

The report (1)

- 1. Describe the problem you want to study. Why is this interesting? What prior knowledge do you have? What do you want to achieve?
- 2. Selection of factors and levels: Which factors do you think are relevant to the problem described above? Which of these factors do you think is active/inert? Do you expect an interaction between some of the factors? Which levels should be used, and why do you think these are reasonable? How can you control that the factors really are at the desired level?
- 3. Selection of response variable: Which response variable will provide information about the problem described above? Are there several response variables of interest? How should the response be measured? What can you say about the accuracy of these measurements?

42 / 45

I don't want to collect data!

- Well, it is possible to instead analyse a observational data set (but talk to the lecturer first),
- or to perform a simulation experiment to investigate properties of the regression model.

Supervision?

 See course page - several possibilities until deadline for hand-in on Tuesday May 2.

45 / 45

DOE Effects

For each By in the award (except po) we define on effect to be

Why? p; gives the change (in y) when ity governors to 1, while Effect; gives the change when X; good from -1 to 1.

This (unformately) nazar that $\frac{\nabla^2}{N}$ Vor (Effect;) = Via (2. $\frac{2}{N}$) = 4. Vor ($\frac{2}{N}$) = $\frac{42}{N}$ Until $\frac{2}{N}$ Until $\frac{2}{N}$ United $\frac{2}{N}$

DOE main effect: 2%, for 14,8,0 Shown in main effects plot



2


PART 4: DOE Effects & Infrance THAY267 L18 2403.2017

Ex: Lime beens 23.

tx: Lime beens 2. .
Write down the regression model with all possible

Interpret by: increase to with one wit >

1



When the two lines in the intraction plot ere perallel => there is no interection effect.

⇒ DOE report (Ex4) ← add explanation for one Westion effect.

3

Significent effects

2) Fit reduced model -> read off:

Curiosity: Selject can be content of from the fall month lay Selject's the Z. Elfects

all Effects not passed oward

3) heath's method.

â

TMA4267 Linear Statistical Models V2017 (L19)

Part 4: Design of Experiments
Blocking
Fractional factorial designs

Mette Langaas

Department of Mathematical Sciences, NTNU

To be lectured: March 28, 2017

DOE workflow

- 1. Set up full factorial design with k factors in R, and
- 2. randomize the runs.
- 3. Perform experiments, and enter data into R.
- 4. Fit a full model (all interactions).
- 5. If you do not have replications, look at Pareto plots and, use this to suggest at reduced model (if possible). Refit the reduced model.
- 6. Assess model fit (residual plots, need transformations?).
- 7. Assess significance.
- 8. Interpret you results (main and interaction plots).

1/23

Genuine run replicates

"When genuine run replicates are made under a given set of experimental conditions, the variation between the associated observations may be used to estimate the standard deviation of the effects. By genuine run replicated we mean that variation between runs made at the same experimental conditions is a reflection of the total variability afflicting runs made at different experimental conditions. This point requires careful consideration." From Box, Hunter, Hunter (1978, 2005): "Statistics for Experimenters", Ch.10.6.

3 / 23

Q: Randomization

Why do you need to randomize the order in which you perform the experiments?

To make the experiments

- A: random.
- ▶ B: robust to external factors
- C: have constant variance.
- D: independent.

Vote at clicker.math.ntnu.no, TMA4267 classroom.

2 / 23

Genuine run replicates

Randomization of run order usually ensures that replicates are genuine. Pilot plant example: each run consists of

- 1. cleaning the reactor
- 2. inserting the appropriate catalyst charge
- 3. running the apparatus at at given temperature and a given feed concentration for 3 hrs to allow the process to settle down at the chosen experimental conditions, and
- 4. combining chemical analyses made on these samples.

A genuine run replicate must involve the taking of all these steps again. In particular, several chemical analyses from a single run would provide only an estimate of analytical variance, usually only a small part of the run-to-run variance. From Box, Hunter, Hunter (1978, 2005): "Statistics for Experimenters", Ch.10.6.

Pilot plant: A, B and C

A=Temperature, B=Concentration, C=Catalyst, Y=vield,

~- I	A reinperature, b—concentration, c—catalyst, r—yield.												
Α	В	C	AB	AC	BC	ABC	Level code	Response					
-	-	-	+	+	+	-	1	60					
+	-	-	-	-	+	+	a	72					
-	+	-	-	+	-	+	b	54					
+	+	-	+	-	-	-	ab	68					
-	-	+	+	-	-	+	с	52					
+	-	+	-	+	-	-	ac	83					
-	+	+	-	-	+	-	bc	45					
+	+	+	+	+	+	+	abc	80					
<i>x</i> ₁	<i>x</i> ₂	<i>X</i> 3	X ₁₂	X ₁₃	X23	X ₁₂₃		у					

5 / 23

Blocking on ABC

C1	C2	C3	C4	CS	O6	C7	CB	C9	C10	C11	
StdOrder	RunOrder	CenterPt	Blocks	Α	В	С	ABC		Y	block effect	
- 1	- 1	- 1	- 1	-1	-1	-1	-1	- 1	60	60	
4	- 4	- 1	- 1	-1	- 1	- 1	-1	7	45	45	
3	3	- 1	1	1	-1	- 1	-1	6	83	83	
2	2	- 1	1	1	- 1	-1	-1	- 4	68	68	
7	7	- 1	2	-1	-1	- 1	- 1	5	52	62	
6	6	- 1	2	-1	- 1	-1	- 1	3	54	64	
- 5	5	- 1	2	1	-1	-1	- 1	2	72	82	
8	8	- 1	2	- 1	- 1	- 1	- 1	8	80	90	

- ► ABC is counfunded with the block effect. We can not separate these two effects from eachother.
- ▶ Suppose all values in block 2 is increased by 10 units.
 - ▶ Then the estimated effect of ABC will increase by 10.
 - ► But all other estimated effects remain unchanged and these are the most important to estimate.

Blocking on ABC

B*C

Block 1 consists of experiments with ABC=-1. Block 2 consists of experiments with ABC=1.

C1	C2	C3	C4	C5	C6	C7	C8	C9	C10	C11	
StdOrder	RunOrder	CenterPt	Blocks	Α	В	С	ABC		Y	block effect	
1	1	1	1	-1	-1	-1	-1	1	60	60	П
4	4	1	1	-1	1	1	-1	7	45	45	
3	3	1	1	1	-1	1	-1	6	83	83	
2	2	1	1	1	1	-1	-1	4	68	68	
7	7	- 1	2	-1	-1	1	1	- 5	52	62	
6	6	- 1	2	-1	1	-1	1	3	54	64	
5	5	1	2	1	-1	-1	1	2	72	82	
8	8	- 1	2	1	1	1	1	8	80	90	

6/23

Original da	ata		Added 10 to all obs in Block 2.				
Factorial Y versus Block A I			Factorial Fit: "block effect" versus Block A B C				
Term	Effect	Coef					
			Term	Effect	Coef		
Constant		64,250	Constant		69,250		
Block		-0,250	Block		-5,250		
A	23,000	11,500	Α	23,000	11,500		
В	-5,000	-2,500	В	-5,000	-2,500		
C	1,500	0,750	C	1,500	0,750		
A*B	1,500	0,750	A*B	1,500	0,750		
A*C	10,000	5,000	A*C	10,000	5,000		

0,000 0,000 B*C

8 / 23

0,000 0,000

2³ with four blocks

We need two generators (columns) to define four blocks: the optimal choice is AB and AC $\,$

▶ Block 1: AB=AC=-1 (- -)

▶ Block 2: AB=-1, AC=1 (- +)

▶ Block 3: AB=1, AC=-1 (+ -)

▶ Block 4: AB=AC=1 (+ +)

Std order	Α	В	С	AB	AC	BC	ABC
1	-	-	-	+	+	+	-
2	+	-	-	-	-	+	+
3	-	+	-	-	+	-	+
4	+	+	-	+	-	-	-
5	-	-	+	+	-	-	+
6	+	-	+	-	+	-	-
7	-	+	+	-	-	+	-
8	+	+	+	+	+	+	+

9 / 23

2³ with AB and AC as generators

- ► Interaction effects AB and AC are confounded with the block effect, since they are the generators.
- ► Their product, AB * AC = A²BC = BC, is also confounded with the block effect (see that BC is constant within each block).
- ▶ Adding h₂ to block 2, h₃ to block 3 and h₄ to block 4 does not change the estimated main effects A, B, or C, and not the interaction effect ABC.
- ▶ However, AB will change with $2 \cdot h_3 + 2 \cdot h_4 2 \cdot h_2$, and we will NOT be able to separate the true AB effect from the block effect.

2³ with AB and AC as generators

Std order	Α	В	С	AB	AC	BC	ABC	Block
2	+	-	-	-	-	+	+	1
7	-	+	+	-	-	+	-	1
3	-	+	-	-	+	-	+	2
6	+	-	+	-	+	-	-	2
4	+	+	-	+	-	-	-	3
5	-	-	+	+	-	-	+	3
1	-	-	-	+	+	+	-	4
8	+	+	+	+	+	+	+	4

10 / 23

How to choose which blocks to be used for blocking?

- Idea: try to leave estimates for main effects and low order interaction unchanged by the blocking.
- ▶ Note: I=AA=BB=CC, where I is a column of 1's.
- ► How NOT to do this:
 - \blacktriangleright Find the blocks for a 2^3 experiment using generators ABC and AC.
 - ▶ The interaction between ABC and AC is ABC*AC=B.
 - ► This means chosing ABC and AC is not a good idea since then we can not trust our estimate of B.

12 / 23

Questions

Should you use a blocking factor in your compulsory project?

Do you understand the difference between blocking and repetition?

13 / 23

Reactor data: standard order

15 / 23


Box, Hunter, Hunter: Reactor example

- ► A=feed rate (liters/min).
- ► B=Catalyst (%).
- ► C=Agitation rate (rpm).
- ▶ D=Temperature (deg C).
- ► E=Concentration (%).
- ► Response= (%) reacted.

Full factorial with $2^5=32$ experiments. From Box, Hunter, Hunter (1978, 2005): "Statistics for Experimenters", Ch.12.2.

14/23

Pareto and Normal plot



Redundancy

- ► The number of runs in a full 2^k factorial design increases geometrically when k is increased.
- ▶ E.g. k = 7 factors gives $2^7 = 128$ runs and we can estimate
 - $(^{7}_{1}) = 7$ main effects
 - $(\frac{7}{2}) = 21$ 2nd order interactions
 - $(\frac{7}{3}) = 35$ 3rd order interactions
 - (7) = 35 4th order interactions
 - $\binom{7}{5} = 21$ 5th order interactions
 - $\binom{7}{6} = 7$ 6th order interactions
 - $(\frac{7}{7}) = 1$ 7th order interactions

17 / 23

Full 2³ factorial experiment

How can we accomodate four factors here?

Std order	Α	В	С	AB	AC	BC	ABC
1	-	-	-	+	+	+	-
2	+	-	-	-	-	+	+
3	-	+	-	-	+	-	+
4	+	+	-	+	-	-	-
5	-	-	+	+	-	-	+
6	+	-	+	-	+	-	-
7	-	+	+	-	-	+	-
8	+	+	+	+	+	+	+

19 / 23

Redundancy (cont.)

- ► There is a hierarchy in absolute magnitude: the main effects tend to be larger than the 2nd order interactions, which tends to be larger than the 3rd order interactions, which ...
- ► At some point higher order interactions tend to become negligible and can be discarded.
- ► If many factors are introduced into a design, it often happens that some have *no* distinguishable effect at all.
- ▶ Fractional factorial designs exploit this redundancy!

18 / 23

Full 2³ factorial experiment - turned into 4-factor experiment

Which effects are confounded?

	Α	В	С	AB	AC	ВС	D=ABC	ABD	ACD	BCD	ABCD
1	-	-	-	+	+	+	-	-	-	-	+
2	+	-	-	-	-	+	+	-	-	+	+
3	-	+	-	-	+	-	+	-	+	-	+
4	+	+	-	+	-	-	-	-	+	+	+
5	-	-	+	+	-	-	+	+	-	-	+
6	+	-	+	-	+	-	-	+	-	+	+
7	-	+	+	-	-	+	-	+	+	-	+
8	+	+	+	+	+	+	+	+	+	+	+

Half fraction of 2⁴

- ▶ The design is called 2_{IV}^{4-1} .
- ▶ D=ABC is called the *generator* for the design.
- ▶ I=ABCD is called the *defining relation* for the design.
- ► The design is said to have *resolution IV*.
- ▶ The *alias structure* defines which effects are confounded:
 - ► A+BCD, B+ACD, C+ABD, D+ABC.
 - ► AB+CD, AC+BD, BC+AD.

21 / 23

What did we learn today?

- Why may experiments need to be performed in blocks? (Batches of raw material, performed on different days, different people performing the experiments.)
- Should we also add a "block" effect if we perform repeated experiments? (Sometimes. If done by different people, or external factors have changed.)

22 / 23

What did we learn today?

Why may experiments need to be performed in blocks?
 (Batches of raw material, performed on different days, different people performing the experiments.)

22 / 23

What did we learn today?

- Why may experiments need to be performed in blocks?
 (Batches of raw material, performed on different days, different people performing the experiments.)
- ► Should we also add a "block" effect if we perform repeated experiments? (Sometimes. If done by different people, or external factors have changed.)
- ► Should then the block effect be a part of the regression model? (In most cases: yes!)

What did we learn today?

- Why may experiments need to be performed in blocks?
 (Batches of raw material, performed on different days, different people performing the experiments.)
- Should we also add a "block" effect if we perform repeated experiments? (Sometimes. If done by different people, or external factors have changed.)
- Should then the block effect be a part of the regression model? (In most cases: yes!)
- Why don't we want to perform a full factorial experiment, but a instead a fractional factorial? (If we have many factors we maybe not need to be able to estimate all possible interactions, and may accept that effects are confounded.)

22 / 23

What did we learn today?

- ▶ What is the easiest way to design a half-fraction of a 2^k factorial experiment? (Perform all the experiments where the highest order interaction =-1 or +1. E.g. for k=4 we may do 16 different experiments, and now we only do the 8 possible experiments where ABCD=+1=defining relation. This is the same as thinking that D=ABC=generator).
- ▶ New words: generator(s), defining relation(s), resolution.

23 / 23

What did we learn today?

▶ What is the easiest way to design a half-fraction of a 2^k factorial experiment? (Perform all the experiments where the highest order interaction =-1 or +1. E.g. for k=4 we may do 16 different experiments, and now we only do the 8 possible experiments where ABCD=+1=defining relation. This is the same as thinking that D=ABC=generator).

23 / 23

What did we learn today?

- ▶ What is the easiest way to design a half-fraction of a 2^k factorial experiment? (Perform all the experiments where the highest order interaction =-1 or +1. E.g. for k=4 we may do 16 different experiments, and now we only do the 8 possible experiments where ABCD=+1=defining relation. This is the same as thinking that D=ABC=generator).
- ▶ New words: generator(s), defining relation(s), resolution.
- Next time: more on interpreting "confounding", interpreting "resolution" and more fractional factorial experiments

Pert 4: DoE

Performing a full 24 factor of expr.

Two important aspects:

- a) The run order is rendown, so that potential external factors one not confused/confounded with experimental factors.
- b) Each experient is a genuine run replicate, that is, reflects the total vanishilly of the experiment.

Blocking

We sold perform a 2s experiment, but howeto use two bookches of new makes of the sound to divide the 8 runs into two groups. When is the book way to do this?

Solution, was the ARC column to define the lolocko, ARC is the block agreement.

ABC=-1; use betch 1
ABC=+1; 2

1

give only run significant effects.

20 in four blocks

To divide the 2^5 =8 runs who 4 block we need two block generates. The best solutions to we Aib and AC as garaceters for the blocks:

Then the block effect will not be confounded with the main effect A,B,C or ABC intradion.
But will be confounded with AB,AC and also
AB.AC. A^BC > BC

Q: What if AEC and EC were to be chosen as black generates $^{\rm R}$

ABC. BC = AOFC== A = blocho vill be conformed with the A effect

3

"The block "Verseble" will be a new regressor replacing the ABC fector.


What would happen if I did not include the olack as a regressor/coverible in the analysis?

SSE will be long.

Example: person as block

Erna continue find user to perform an 25 experiment trighter, and to get to obs. they will both conduct the same physical 2° experiments.

Should then a consonel telling who did actorphisms be added to the respection model?



If the above figur gives a correct picture of the experiment NOT including a person coverace will make SSE voy legge, and will threfore

2

Fractional factorial designs

Observation: when the window of factor (k) is large, it may not be optional to perform a full 24 factoral duringly due to the possible redundency of the during.

higher-order interections than to be smaller than lower order interections

Solution: only purform a fraction of the full design!

Now: move in the opposite direct on to solve this.

We have a full 2^{α} factorial durign with factor A B_1 G but we also wrent to have a new factor D in the experiment.

Possible solution: let the AGC column define the favelo of factor O.

1) D= ABC is called the "generator" of the danger.

2) I = D.D = D. ABC = ABCD is called object the defining relation of the design.

4

- 3) The number of letters (lenth) of the (shoriest) defining relation is called the "resolution of the design", and is landed by Roman nunarels. Here: III
- 4) Finally: this is called a half fraction of a 24 design

Not ATION

2 design with I = ABCD as delians relation and D= ADC es generater.

may use Lenth's netword to assess significance

with 4 factors then ere:

- 4 = (1) main effect A, O, C, O
- 6 = (4) two-vay invecting As, AC., Co
- 4 = (4) three-way interestors ABC ACO, BCO
- 1 = (4) four-way nurection ABCD

4+6+4+1= 15 possible effect (+1 insucapt) =

2) 2-way:

AB: AB: T = AB. ABCD - CD las = AB+CD AC = AD:

RAC = AC+ BD Las = AD +BC

- 3) 3-way: already done a main effects
- 4) I. ABCO is confounded with salvapt.

7

a: what over we entirese? The "alian-stucture" defines which effects are confounded. Obvious Since D: ABC, then Dand AGC ex. confounded. D may actually lee D+ASC Method: We want to find of any effects are confounded with A. We multiply A with the defining relation, I=ABCO. 1) Main effects: A= A.I = A . A6CD = A 6CD = 6CD that is A and OCD column are agral B= B.I = B. ABCP= ACD C = C.T = C. ABCD = ABD D = D.T = D. AGCD = AGC All main affects are confounded with Sway intractions LA = A + 600 we think that we solimate A, but we actually estimate A+BCD, BUT if 8-way nurediences and => On.

TMA4267 Linear Statistical Models V2017 (L20)

Part 4: Design of Experiments Fractional factorial designs Quiz with Kahoot!

Mette Langaas

Department of Mathematical Sciences, NTNU

To be lectured: March 30, 2017

What did we learn last lession?

- Why don't we want to perform a full factorial experiment, but a instead a fractional factorial? (If we have many factors we maybe not need to be able to estimate all possible interactions, and may accept that effects are confounded.)
- ▶ What is the easiest way to design a half-fraction of a 2^k factorial experiment? (Perform all the experiments where the highest order interaction =-1 or +1. E.g. for k=4 we may do 16 different experiments, and now we only do the 8 possible experiments where ABCD=+1=defining relation. This is the same as thinking that D=ABC=generator).
- ► New words:
 - ▶ generator(s)=how to generate the design,
 - defining relation(s), found from the generators,
 - ► resolution=length of shortest defining relation,
 - alias structure=confounding pattern, found by multiplying each effect of interest with the defining relation.
- ► Today: more on interpreting "confounding", interpreting "resolution" and more fractional factorial experiments

1/19

Half fraction with reactor example

- ▶ Instead of running a full factorial with $2^5 = 32$ experiments,
- ▶ we suggest running a half-fraction.
- We choose I = ABCDE as the defining relation.

3/19

Box, Hunter, Hunter: Reactor example

- ► A=feed rate (liters/min).
- ► B=Catalyst (%).
- ► C=Agitation rate (rpm).
- ► D=Temperature (deg C).
- ► E=Concentration (%).
- ► Response= (%) reacted.

Full factorial with $2^5 = 32$ experiments. From Box, Hunter, Hunter (1978, 2005): "Statistics for Experimenters", Ch.12.2.

2/19

Reactor data: answer in groups

- ► Which of the 32 experiments should be performed when *I* = *ABCDE* is the defining relation? What is then the generator?
- ▶ What is the resolution for this design?
- ▶ Write down the aliasing pattern.

Resolution

A design is said to be of resolution R if no p-factor effect is aliased with an effect containing less than R-p factors.

A design of resolution

- III does not confound main effects with one another, but does confound main effects with two-factor interactions.
- IV does not confound main effects and two-factor interactions, but does confound two-factor interactions with other two-factor interactions.
- V does not confound main effects and two-factor interactions with each other, but does confound two-factor interactions with three-factor interactions and so on.

In general the resolution of a two-level factional design is the length of the shortest word in the defining relation.

5/19

7/19

Interpretation of confounding: example

Suppose there are three factors, A, B, C, for which we know the true effects and interaction effects:

A = 8

B = 20

C = 2

AB = 4

AD - 4

AC = 2

BC = 6

ABC = 4

Also is known that average response is 70.

Half fraction with reactor example: generator and defining relation

- ▶ Instead of running a full factorial with $2^5 = 32$ experiments,
- ▶ we suggest running a half-fraction.
- ▶ We choose I = ABCDE as the defining relation.
- ► Alternative thinking:
 - ► Construct a full 2⁴ design for A, B, C and D.
 - ► The column of signs for the ABCD interaction is written and used to define the levels for factor E.
 - ► This means E = ABCD is the generator for the design, and I = ABCDE is the defining relation.

R-code on course www-page.

6/19

True regression model

The corresponding regression model is:

$$y = \beta_0 + \beta_1 z_1 + \beta_2 z_2 + \beta_3 z_3 + \beta_{12} z_{12} + \beta_{13} z_{13} + \beta_{23} z_{23} + \beta_{123} z_{123} + \epsilon$$

where $z_{12}=z_1z_2$, $z_{13}=z_1z_3$, $z_{23}=z_2z_3$, $z_{123}=z_1z_2z_3$, and where the coefficients β are half the corresponding effects, while $\beta_0=70$. The regression model is hence

$$y = 70 + 4z_1 + 10z_2 + z_3 + 2z_{12} + z_{13} + 3z_{23} + 2z_{123} + \epsilon$$

In the following we shall also for simplicity assume that the errors ϵ are 0. This makes it possible to compute the responses for any experiment for which the levels of A, B, C are specified.

Confounding example (cont.)

Assume now that a 2^{3-1} experiment is performed, with generator C=AB. And responses are computed using the true regression model (check!).

St. order		Α	В	C=AB	AB	AC	BC	ABC	у
1	+	-	-	+	+	-	-	+	57
2	+	+	-	-	-	-	+	+	65
3	+	-	+	-	-	+	-	+	73
4	+	+	+	+	+	+	+	+	93
	Const.	z_1	<i>z</i> ₂	<i>Z</i> 3	Z ₁₂	z ₁₃	Z ₂₃	Z ₁₂₃	
Coeff.	70	4	10	1	2	1	3	2	

9/19

Confounding example (cont.)

 Suppose now that we try to compute the main effect of A from our data. Apparently this will be

$$\ell_A = \frac{65+93}{2} - \frac{57+73}{2} = 79-65 = 14$$

which is also found as twice the coefficient before z_1 in the regression model above.

 Similarly, the apparent interaction effect of B and C would be computed as

$$\ell_{BC} = \frac{-57 + 65 - 73 + 93}{2} = 14$$

The truth (which is known to us) is, however, that A=8 and BC=6, so that it is the sum of A and BC which is 14.

11 / 19

Confounding example (cont.)

It is now seen that in all of these 4 experiments are

Const. =
$$z_{123}$$

 $z_1 = z_{23}$
 $z_2 = z_{13}$
 $z_3 = z_{12}$

so for the performed experiment we may as well write the model as

$$y = (\beta_0 + \beta_{123}) + (\beta_1 + \beta_{23})z_1 + (\beta_2 + \beta_{13})z_2 + (\beta_3 + \beta_{12})z_3$$

Using that we know the values of the coefficients, the true model for the data is thus

$$y = (70+2) + (4+3)z_1 + (10+1)z_2 + (1+2)z_3$$

= 72 + 7z₁ + 11z₂ + 3z₃

10 / 19

This is what is meant by saying that the main effect of A and the interaction effect between B and C are *confounded* (mixed). The confounded effects are listed in R as the *alias structure*.

Factorial Fit: y versus A; B; C

Estimated Effects and Coefficients for y (coded units)

Term	Effect	Coef
Constant		72,000
A	14,000	7,000
В	22,000	11,000
C	6,000	3,000

Alias Structure

I + A*B*C A + B*C B + A*C C + A*B

The bicycle example

TABLE 12.5. An eight-run experimental design for studying how time to cycle up a hill is affected by seven variables (I = 124, I = 135, I = 236, I = 1237).

run	seat up/down 1	dynamo off/on 2	handlebars up/down 3	gear low/medium 4 12	raincoat on/off 5	breakfast yes/no 6 23	tires hard/soft 7 123	time to climb hill (sec) y
!		_	_	+	+	+	-	69
2	+	-	-	_	_	+	+	52
3	_	+	-	-	+	-	+	60
4	+	+	_	+	_	-	-	83
5	_	_	+	+	_	_	+	71
6	+	-	+	_	+	-	_	50
7	_	+	+	-	_	+	-	59
8	+	+	+	+	+	+	+	88

From Box, Hunter, Hunter (1978, 2005): "Statistics for Experimenters", Ch.12.25

13 / 19

Using FrF2 in R, see file L20.R

15 / 19

The bicycle example

- ▶ Set up a full factorial design in the three variables A, B, C.
- ▶ Use the generators: D=AB, E=AC, F=BC, G=ABC.
- ▶ Defining relations: I=ABD=ACE=BCF=ABCG.
- ▶ The design is of resolution III.
- ▶ It is a 1/16 fraction of the full 2^7 , and thus called 2_{III}^{7-4} .
- ► A design where every available contrast is associated with a factor is called a *saturated design*.

14 / 19

Exam question on fractional factorials (K2014)

In a pilot study with four factors A, B, C and D, the 8 experiments listed below were run.

1
-1
-1
1
1
-1
-1
1

What type of experiment is this?


What is the generator and the defining relation for the experiment?

What is the resolution of the experiment?

Write down the alias structure of the experiment.

Not covered: Response Surface Methods

Dates back to the 1950s, with popular book by Box and Draper.



- The method performes sequential optimization, and can deal with several responses simultaneously.
- Central Composite Designs (CCD) and Box-Behnken Designs are two popular methods.
- ➤ John Tyssedal supervises 5th year project and master thesis in DOE.

https://onlinecourses.science.psu.edu/stat503/node/57

17 / 19

Summing up with Kahoot! quiz




kahoot.it

19 / 19

Final word about the DOE Compulsory Exercise 4

- ► If you want to have 4 factors and perform 16 runs see R-code named https://www.math.ntnu.no/emner/TMA4267/ 2017v/RscriptDOEtreadmill.R
- ▶ If you want to have 3 factors, but need a block effect look at this code https://www.math.ntnu.no/emner/TMA4267/2017v/D0E2in3withrepl.R, because it is best to code the block with effect coding FrFr use treatment coding and then we don't have orthogonal columns and everything becomes difficult...



Interpretation of confunds

- 1) y= 70 + 451+ 102+ 12+22+320+
- 2) C=AB, A=BC, B=AC, AOC=I
- 3) y=72+72,+1122+323

We think we exhaut A, but really exhaust A+BC=14.

Finally: bicycle example Hove 7 factors and perform 8 expersants \$\$2^{-4}\$ design. Thin: fell factoration A, a, c and add generation De

see L20.R for e-coax.

2

$$Y = X\beta + \varepsilon$$
, with $\varepsilon \sim N_n(0, \sigma^2 I)$
 $\widehat{Effect}_j = 2 \cdot \frac{1}{n} \sum_{i=1}^n x_{ij} Y_i$.

 $Var(Effect_j)$ equals

 σ^2


 $\frac{1}{n}\sigma^2$

 $c \frac{2}{n} \sigma^2$

 $D \frac{4}{n}\sigma^2$


Which of the following is NOT correct for a 2^k full factorial design matrix X?

- **A** X only contains the numbers -1 and 1.
- B The sum of each column equals 1.
- \mathbf{c} The columns of \mathbf{X} are orthogonal.
- $\mathbf{D} \mathbf{X}^T \mathbf{X}$ is a diagonal matrix.



This plot is called

- A Main effects plot
- **B** Interaction effects plot
- c Pareto plot
- Normal plot




Set up a full factorial design in the three variables A, B, C, and use generators: D=AB, E=AC, F=BC, G=ABC. What do you get?

 A 2_{III}^{7-4}

 $^{\rm B}$ 2^{7-3}_{IV}

 $c 2^{7-4}_{IV}$

 $^{\circ}$ 2_{III}^{7-3}



For a design is of resolution III:

- A Main effects are confounded with each other.
- B Main effects are confounded with 2-way interactions.
- Main effects are confounded with 3-way interactions.
- D Main effects are confounded with 4-way interactions.

Correct?

Are you sure you want to read the correct answers? Maybe try first?

Answers

Correct: BDCDAAB