

Slides Week 38

Some properties of Statistics

X_1, \dots, X_n are $N(\mu, \sigma^2)$

$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ and $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ are independent

$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$, $\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$

T-statistic: $\frac{\bar{X} - \mu}{\frac{S}{\sqrt{n}}}$, In general $T_p = \frac{N(0,1)}{\sqrt{\frac{\chi^2(p)}{p}}}$

$$\text{Var}[T_p] = \frac{p}{p-2}$$

$$F_{p,q} \text{ statistic} = \frac{\frac{\chi^2(p)}{p}}{\frac{\chi^2(q)}{q}}$$

$$V \sim \chi^2(q) \Leftrightarrow V \sim \Gamma\left(\frac{q}{2}, 2\right)$$

$$E(V^{-k}) = \frac{1}{\Gamma(q/2)2^{q/2}} \int_0^{\infty} v^{q/2-k-1} e^{-v/2} dv = \frac{\Gamma(\frac{q}{2} - k)}{\Gamma(\frac{q}{2})2^k},$$

$$E[F] = \frac{q}{q-2}$$

$$\text{Var}[F] = \frac{2q^2(q+p-2)}{p(q-2)^2(q-4)}$$